



**Urban EC and OC Trends**  
**Using CSN, IMPROVE, SEARCH EC & OC,**  
**“BC” & OCMmb via FRM PM2.5**  
*Findings and Issues*

Neil Frank US EPA / OAQPS  
For Presentation at IMPROVE Carbon Workshop  
Columbia Gorge, Washington  
Oct 27, 2010

# Outline

## Background

### Part 1: Urban EC trends

- EPA's CSN, 2002-mid 2010
- CSN inter-comparisons
  - IMPROVE TOR 3000N EC v. NIOSH TOT MetOne EC
  - SEARCH vs CSN (Birmingham)
- Longer-term trends
  - Washington DC (1990-2009)
  - BC from Boston and S.F.

### Part 2: Urban OC Trends

- CSN OC vs SANDWICHed OCMmb, 2002-mid 2010
- TC vs TC & OC:EC ratios

# Background

## CSN Carbon Measurements (~200 urban sites)

- **Until 2007 -- NIOSH-like TOT protocol,**
  - mostly using MetOne sampler
- **Starting May 2007, Switched to IMPROVE TOR protocol with URG 3000N**

## Carbon Measurements Are Also Affected by Sampler & Filter Face Velocity

- **MetOne - 6.7 Lpm, 47 mm quartz**
- **FRM - 16.7 Lpm , 47mm Teflon**
- **URG 3000N - 22.8 Lpm, 25mm quartz (highest FV)**
  - ~12x MetOne,
  - ~ 3x PCM3,
  - ~1.3x FRM (w. McDade et al [2009] filter screen adjustment)
- **SEARCH PCM3 (w IMPROVE TOA) - 16.7 Lpm , 37mm quartz and denuder**

# *“Conventional” Thinking*

- **EC TOR > EC TOT**
- **IMPROVE TOR EC > NIOSH TOT EC**
- **OC is primarily affected by positive sampling artifact**
- **EC does not have sampling artifacts**


*Some modifications may be needed for CSN !*



# CSN EC Trends, 2002-mid 2010

# All CSN sites with NIOSH-like protocol


Distribution of monthly mean **EC** among all sites  
(Individual sites may not display strong seasonality)



PLOT    NIOSH\_EC    NIOSH\_EC    NIOSH\_EC    NIOSH\_EC


# 74 selected for Trends (most complete CSN carbon sites)

EC Concentration,  $\mu\text{g}/\text{m}^3$



- 53 sites which switched in May 2007
- IMPROVE protocol
  - URG 3000N (IMPROVE like sampler)

EC Concentration,  $\mu\text{g}/\text{m}^3$




PLOT Avg\_ec\_3DDDN Avg\_ec\_3DDDN Avg\_ec\_3DDDN Avg\_ec\_3DDDN

## TO REMOVE CHANGING SITES and Completeness:







- 18 Selected Sites with >50 measurements per year, 2002-2009

*Is there a discontinuity in 2007?*

*Did the trend change direction because of the monitoring change?*




# 18 sites 3000N IMPROVE (site specific trends)








EC declined since 2007 at most sites


# 56 MetOne sites w. NIOSH



# 56 MetOne sites w. NIOSH

The decline after 2007 appears to be nationwide






# TO ESTIMATE THE APPARENT DISCONTINUITY IN 2007

# Step 1

## 2 Month Collocation (May-June 2007)

- 3000N = 14% higher than CSN MetOne NIOSH




## Step 2

### 12-month Carbon Inter-Comparison of 11 CSN sites

NIOSH OC/EC w. MetOne sampler

vs. IMPROVE OC/EC w. URG 3000N sampler



Bronx and Queens, NY;

Atlanta GA,

Birmingham AL,

Detroit MI;

Cleveland OH;

Chicago IL,

Denver CO,


LA (Rubidoux) CA;

Sacramento CA \*

Seattle WA \*


\* woodsmoke sites

May 1009-April 2010, (not presented chronologically)




ratio on days then CSN NIOSH > 0.1, all days shown for trend lines

# Many Sites show 3000N EC > MetOne EC Summer only Pattern



# 24-hr EC measurements (897, May 2009-April 2010)

EC 3000N




Cool Months - Oct-Mar

Warm Months - Apr-Sep

CSN NIOSH EC is low in part  
due to low warm season EC values

NIOSH EC

# 24-hr EC measurements (897, May 2009-April 2010)



**Without NIOSH EC < 0.05 ug/m3**

**TOR= 0.81 \* TOT (Oct-Mar)**


**TOR= 1.14 \* TOT (Apr-Sep)**

PLOT    ●-●-● w\_ec\_3000N    ◻-◻-◻ S\_ec\_3000N


Regression Equation:  
w\_ec\_3000N = 0 + 0.818246 \* NIOSH\_EC  
S\_ec\_3000N = 0 + 1.146079 \* NIOSH\_EC

w\_ec\_3000N

# MONTHLY AVERAGE EC (May 2009-April 2010)




Regression Equation:  
w\_ec\_3000N = 0 + 0.860958\*niosh\_ec  
s\_ec\_3000N = 0 + 1.2828\*niosh\_ec




Regression Equation:  
 $w\_ec\_3000N = 0 + 0.857535 * niosh\_ec$   
 $s\_ec\_3000N = 0 + 1.192341 * niosh\_ec$

*During 2009-2010,  
 there appears to be seasonal relationship* 21

# EC Inter-Comparison, by Month (11 CSN sites, May 2009-Apr 2010)








## **Birmingham EC (2000-2010)**

### **3 collocated METHODS**

**What can we learn from comparison to SEARCH's EC**

# SEARCH EC in Birmingham confirm EC decrease but relative CSN to SEARCH concentrations have changed






New 2009 agreement suggests NIOSH EC is too high




New disagreement suggests 3000N is too low


# 2009 EC in Birmingham

- 3000N TOR is consistently lower than CSN NIOSH TOT



Also, 3000N TOR is lower than SEARCH EC TOR






**National CSN EC trend**  
**Tentatively “HARMONIZED” to account for**  
**network change**

# EC 3000N is adjusted to be NIOSH-like

Using simple 2009-2010 relationship:


(i.e. 20 % higher in the summer, 10-20% lower in the winter)



# EC NIOSH is adjusted to be 3000N-like

Using simple 2009-2010 relationship:


(i.e. 20 % lower in the summer, 10-20% higher in the winter)







# **Urban EC & BC before 2002**

**What can we say about longer-term urban EC trends**








# Aethalometer BC

# BC via COHs

## Boston, 2000-2010



## San Francisco (1965-2005)



BC was estimated using Coefficient of Haze measurements which are shown to be highly correlated with optical BC  
 Source: Kirchstetter et al., 2008.

Relatively flat since 2004




## Part 2: OC trends


# 18 Selected Sites with >50 measurements per year, 2002-2009

*Note the differential concentration distribution and possible trend discontinuity*

- Using nominal blank adjustments
  - 1-1.5  $\mu\text{g}/\text{m}^3$  for CSN NIOSH with MetOne
  - 0.4  $\mu\text{g}/\text{m}^3$  for CSN 3000N




# Overlay 18 sites with SANDWICHed OCMmb




# Birmingham

OC and OCMmb, monthly avgs.




$$\text{OCMmb} = a + k \cdot (\text{OC}') \quad \text{OC}' \text{ includes nominal blank corrections of } 1 \text{ and } 0.4 \mu\text{g}/\text{m}^3$$




56 Selected Sites with >50 measurements per year, 2002-2009  
(did not switch to 3000N in 2007)

OC Concentration,  $\mu\text{g}/\text{m}^3$


**OC\_NIOSH (56 sites)**



PLOT Avg\_oC avg\_oc3 Avg\_oC avg\_oc3 Avg\_oC avg\_oc3



# Birmingham, Shows Declines in EC and OC (SEARCH and CSN 3000N)



**Randy Dillard (Jeff Co. AL) is checking for possible changes to CSN instrumentation**

SEARCH Carbon Protocol:

IMPROVE TOR by DRI

Sampler: 16.7 LPM, 37mm Pallflex® Tissuquartz, Denuder, backupfilter


OC (blank corrected, using quarterly average values) = OCQF + OCQBQ - 2OCbQF

backup


field blank



# OC:EC ratios and Sampling Artifacts



**Urban OC:EC ratio**  
**Is the true urban ambient ratio somewhere in between?**




Distrn using Monthly average values, without NIOSH <0.1

43  
 With nominal blank adjustments

Ratios are not the same because TC is not the same  
different CSN samplers, face velocity and artifacts- positive (and negative?)

Low flow rate sampler collects relative more OC  
(and high flow rate sampler collects relatively more)




Regression Equation:  
 $NIOSH\_TC = 0.880438 + 1.092001 * TC\_3000N$

includes nominal values of 1 and 0.4

Note: Chow et al. (2009)  
**CSN NIOSH = 1.65 + 1.08 IMPROVE (PHOE1)**  
(without blank corrections)

# Relative Difference is Seasonal (Additional 2-3 ug/m<sup>3</sup> NIOSH TC)


Represents estimated CSN NIOSH positive artifact and/or lost 3000N semi-volatile OC




## As per Schichtel et.al.,

### •Relating CSN to IMPROVE data:

- $[EC]^{CSN-URG} = [EC]^{IMP}$
- $[OC]^{CSN-URG} = [OC]^{IMP} + A^{CSN-URG}$
- $[TC]^{CSN-MetOne} = [EC]^{IMP} + (1+B^{IMP*})[OC]^{IMP} + A^{CSN-MetOne}$
- $[EC]^{CSN-MetOne} = (1-g)*[EC]^{IMP}$ ; multiplicative bias between EC by TOT and TOR




MetOne OC = A + (1+B) 3000N → A = MetOne positive artifact  
 B = 3000N negative artifact



11 collocated CSN sites (2009-2010), by month - without any OC adjustments

MetOne OC = A + (1+B) 3000N → A = MetOne positive artifact  
B = 3000N negative artifact





# Conclusions/Questions

## Part 1 (EC Trends)

- Urban trends appear level 02-07, followed by 07-09 decline
- Evidence that urban declined during the 1990's thru early 2000's.
- Why do CSN EC agree?
  - Due to sampling artifacts? ...pyrolysis?... and/or different 2009 -10 conditions?
  - Would this relationship hold prior to 2009? Analyze archived filters?
  - e.g. Why does Birmingham behave differently (check samplers....)

## Part 2 (OC Trends, OC:EC)

- What are the CSN sampling artifacts?
  - MetOne -- positive artifact ( $> >1 \text{ ug/m}^3$ ), and partly responsible for high EC (pyrolyzed carbon?)
  - URG3000N -- negative? , and partly responsible for low OC:EC ratio?
- Implications ?...Sampling study needed to estimate negative artifact?

