

LIST OF TABLES

Chapter		Page
	OVERVIEW AND SUMMARY	
	Table S.1 IMPROVE and NPS/IMPROVE protocol sites according to region.	S-3
1	INTRODUCTION	
	Table 1.1 IMPROVE and NPS/IMPROVE protocol sites according to region.	1-6
2	MONITORING METHODOLOGIES	
	Table 2.1 Specifications of the DRI Thermal/Optical Reflectance Carbon Analyzer.	2-10
	Table 2.2 Precisions for average concentration of measured and composite variables.	2-11
	Table 2.3 Transmissometer parameters.	2-14
	Table 2.4 Calculated U_T , Shenandoah summer 1991 study.	2-17
4	VALIDATION	
	Table 4.1 Means and standard errors of sulfur by UCD PIXE in ng/m^3 , for samples with and without a denuder.	4-3
	Table 4.2 $H=b_0+b_1S+b_2$ OM regressions.	4-7
5	SPATIAL AND SEASONAL DISTRIBUTION OF AEROSOL CONCENTRATION	
	Table 5.1 Measured fine and coarse aerosol concentrations (in $\mu\text{g/m}^3$) for the 19 regions in the IMPROVE network.	5-3
	Table 5.2 Measured aerosol mass budgets (in percent) for the 19 regions in the IMPROVE network.	5-6
6	SPATIAL AND SEASONAL DISTRIBUTION OF RECONSTRUCTED LIGHT EXTINCTION	
	Table 6.1 Parameters of the best-fit quadratic equation relating the relative humidity light extinction correction factors (F_T and F_H) to average site relative humidity.	6-8
	Table 6.2 Seasonal and annual averages, averaged over the three-year period, of reconstructed total light extinction coefficient (Mm^{-1}) for the 19 regions in the IMPROVE network.	6-15
	Table 6.3 Seasonal and annual averages, averaged over the three-year period, of reconstructed aerosol light extinction coefficient (Mm^{-1}) for the 19 regions in the IMPROVE network.	6-18
	Table 6.4 Seasonal and annual averages, averaged over the three-year period, of percentage contributions to the reconstructed	6-21

aerosol light extinction coefficient (light extinction budget)
for the 19 regions in the IMPROVE network.

7 MEASURED LIGHT EXTINCTION

Table 7.1	Comparison of measured and reconstructed light extinction coefficient (Mm^{-1}) averaged over the three-year period.	7-4
Table 7.2	Comparison of measured and reconstructed standard visual range (SVR) by region for every season, based on averages of the daily median extinction.	7-6