


Appendix G. Supplemental Analysis for Chapter 9

G.1. REGIONAL HAZE RULE IMPROVE PROGRESS TRACKING SITE DATA RESULTS BY STATE


Baseline (2000-2004) and period 1 (2005-2009) 20% worst and 20% best visibility estimates and natural condition targets for tracking progress under the RHR are presented in this appendix for each of 107 of the 110 regional haze rule tracking sites. Contributions from individual species are also presented. The sites are organized by state so some sites are repeated if they border more than one state. Detailed composition data for every site can be found on the VIEWS website (<http://views.cira.colostate.edu/web/Composition/>).

State	Page
Alabama	G-2
Alaska	G-4
Arizona	G-10
Arkansas	G-26
California	G-30
Colorado	G-64
Florida	G-76
Georgia	G-82
Hawaii	G-86
Idaho	G-90
Kentucky	G-100
Maine	G-102
Michigan	G-106
Minnesota	G-110
Missouri	G-114
Montana	G-118
Nevada	G-134
New Hampshire	G-136
New Jersey	G-138
New Mexico	G-140
North Carolina	G-156
North Dakota	G-164
Oklahoma	G-168
Oregon	G-170
South Carolina	G-182
South Dakota	G-184
Tennessee	G-188
Texas	G-190
Utah	G-194
Vermont	G-200
Virginia	G-202
Virgin Islands	G-206
Washington	G-208
West Virginia	G-220
Wyoming	G-222




Sipsey - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	29.0	28.3	11.0		29.2	28.7	29.4	28.8	31.5	29.7	30.0	26.8	23.6
Sulfate Bext	133.0	129.3	3.3		134.5	130.0	142.2	125.5	196.1	155.5	140.3	95.6	58.8
Nitrate Bext	8.5	7.2	1.0		6.2	10.1	9.6	8.3	3.9	6.6	3.2	9.0	13.5
POM Bext	24.0	24.5	10.8		23.7	23.3	21.5	27.4	20.8	18.7	41.1	25.4	16.7
EC Bext	6.5	6.4	0.3		6.7	5.9	6.2	7.1	8.1	5.7	7.1	5.8	5.2
Soil Bext	0.8	0.8	0.8		0.6	1.0	0.8	0.9	0.7	1.0	1.0	0.9	0.7
Coarse Mass Bext	3.2	3.1	2.4		4.6	3.1	2.4	2.7	3.1	3.0	3.8	2.9	2.5
Sea Salt Bext	0.3	0.2	0.7		0.0	0.0	0.3	0.9	0.1	0.2	0.2	0.1	0.1




Sipsey - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.6	14.1	5.0	14.4	16.4	16.3	15.2	15.7	15.1	14.3	13.3	11.8	
Sulfate Bext	18.5	15.5	0.9	16.3	20.3	20.4	17.0	19.2	18.6	16.5	12.8	10.4	
Nitrate Bext	7.1	4.9	0.9	6.9	8.0	7.6	5.7	6.5	5.9	4.5	4.9	2.8	
POM Bext	6.5	5.5	2.4	5.0	6.9	6.9	7.0	6.3	5.6	5.6	5.5	4.7	
EC Bext	2.6	2.4	0.1	2.3	2.7	2.6	2.8	3.4	2.5	2.3	2.1	1.7	
Soil Bext	0.4	0.3	0.2	0.3	0.5	0.5	0.4	0.4	0.2	0.3	0.3	0.3	
Coarse Mass Bext	2.1	1.8	1.0	1.8	2.3	2.3	2.1	1.8	1.7	2.2	1.6	1.7	
Sea Salt Bext	0.1	0.3	0.1	0.0	0.0	0.0	0.4	0.2	0.2	0.4	0.3	0.1	




Denali - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	9.9	10.6	7.3	10.6	9.1	10.2	9.9	9.4	10.4	9.8	8.8	9.3	14.7
Sulfate Bext	4.9	7.8	1.1	4.6	4.6	6.0	5.4	3.6	4.5	8.5	5.6	7.6	13.1
Nitrate Bext	0.6	0.7	0.6	0.7	0.6	0.4	0.5	0.8	0.6	0.7	0.9	0.8	0.7
POM Bext	10.8	10.7	7.3	23.5	7.1	8.7	7.0	7.9	12.5	3.6	3.9	3.3	30.4
EC Bext	1.6	1.3	0.5	2.3	1.3	1.4	1.6	1.3	1.7	0.9	1.1	0.8	2.1
Soil Bext	0.3	0.3	0.3	0.2	0.5	0.5	0.2	0.2	0.2	0.4	0.2	0.3	0.4
Coarse Mass Bext	1.4	1.2	1.4	1.2	1.8	1.6	1.0	1.3	1.2	1.2	0.9	1.3	1.3
Sea Salt Bext	0.4	0.8	0.6	0.5	0.3	0.0	0.5	0.9	0.7	1.0	0.8	1.0	0.6




Denali - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.4	2.4	1.8	2.7	2.5	2.3	2.2	2.4	2.1	2.5	2.5	2.5	2.5
Sulfate Bext	0.8	0.9	0.2	0.8	1.0	0.8	0.7	0.8	0.5	1.0	0.9	0.9	1.0
Nitrate Bext	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.1
POM Bext	0.3	0.3	0.2	0.5	0.3	0.2	0.3	0.3	0.3	0.3	0.4	0.4	0.3
EC Bext	0.2	0.1	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1
Soil Bext	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.2
Sea Salt Bext	0.1	0.1	0.0	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1




Simeonof - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.6	18.6	15.6		18.8	18.2	18.6	17.4	18.7	20.0	17.7	19.2	
Sulfate Bext	15.3	22.0	1.8		15.2	16.0	14.7	17.5	22.5	23.5	19.0	27.4	
Nitrate Bext	1.9	1.9	1.7		1.8	2.1	1.9	1.8	2.2	1.8	1.9	2.0	
POM Bext	4.6	1.2	2.6		7.5	3.7	2.5	1.8	1.7	0.5	1.3	0.9	
EC Bext	1.9	0.8	0.2		3.8	0.9	1.1	0.9	1.5	0.4	0.4	0.8	
Soil Bext	0.2	0.2	0.2		0.2	0.2	0.2	0.1	0.4	0.2	0.1	0.2	
Coarse Mass Bext	4.4	5.2	3.2		4.4	4.3	4.5	4.5	6.2	6.4	4.4	4.4	
Sea Salt Bext	25.2	23.7	27.5		22.3	24.1	29.0	20.3	21.8	31.6	21.1	23.9	




Simeonof - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.6	8.0	5.3			7.8	6.8	8.3	7.8	8.3	8.1	7.5	8.3
Sulfate Bext	3.7	4.2	0.8			4.3	2.6	4.3	3.6	4.8	4.0	3.8	4.6
Nitrate Bext	0.5	0.5	0.6			0.5	0.4	0.7	0.5	0.6	0.5	0.4	0.5
POM Bext	0.6	0.3	0.7			0.7	0.4	0.7	0.3	0.2	0.2	0.5	0.3
EC Bext	0.4	0.5	0.2			0.4	0.5	0.4	0.4	0.7	0.6	0.4	0.6
Soil Bext	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Coarse Mass Bext	1.1	1.2	0.7			1.1	1.1	1.1	1.1	1.3	1.2	1.1	1.1
Sea Salt Bext	3.2	3.7	2.0			2.9	2.9	3.8	4.1	3.5	4.0	2.9	4.1




Tuxedni - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.1	13.5	11.3			15.5	12.6	14.2	14.2	13.9	12.6	12.9	14.0
Sulfate Bext	8.7	13.0	1.8			8.7	9.8	7.7	10.7	16.2	10.3	12.0	15.9
Nitrate Bext	1.8	1.3	1.6			2.7	1.3	1.3	1.4	1.3	1.3	1.3	1.1
POM Bext	8.9	4.0	5.5			12.3	5.5	8.8	6.5	2.9	4.9	3.3	2.6
EC Bext	1.2	0.9	0.3			1.1	1.0	1.4	1.4	1.0	0.9	0.6	0.5
Soil Bext	0.1	0.4	0.1			0.2	0.1	0.1	0.1	1.5	0.2	0.2	0.3
Coarse Mass Bext	2.5	2.9	2.5			2.9	2.4	2.2	2.8	3.6	2.3	1.7	3.9
Sea Salt Bext	8.2	5.9	8.9			11.3	4.0	9.4	8.7	3.7	5.1	7.0	5.1




Tuxedni - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.0	4.1	3.1			4.2	3.8	4.0	3.9	4.4	3.6	4.1	4.3
Sulfate Bext	1.1	1.4	0.3			1.1	1.0	1.1	1.2	1.4	1.1	1.4	1.7
Nitrate Bext	0.4	0.3	0.4			0.3	0.4	0.4	0.3	0.3	0.2	0.2	0.3
POM Bext	0.4	0.3	0.3			0.5	0.3	0.3	0.3	0.4	0.2	0.3	0.1
EC Bext	0.2	0.1	0.1			0.1	0.2	0.3	0.1	0.1	0.0	0.1	0.1
Soil Bext	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Coarse Mass Bext	0.3	0.5	0.3			0.4	0.3	0.3	0.4	0.6	0.3	0.4	0.5
Sea Salt Bext	0.5	0.6	0.4			0.7	0.4	0.5	0.5	0.7	0.5	0.7	0.8




Chiricahua - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.4	12.2	7.2	14.0	12.5	14.5	14.9	11.2	12.6	13.4	12.3	12.1	10.7
Sulfate Bext	8.1	9.1	0.6	8.7	8.5	7.8	8.6	7.0	10.6	8.3	10.7	9.0	7.0
Nitrate Bext	1.3	1.3	0.7	1.1	0.9	1.9	1.5	1.4	1.0	1.7	1.1	1.1	1.4
POM Bext	7.3	4.0	3.9	8.5	4.5	9.3	9.8	4.3	5.4	4.5	3.8	4.1	2.5
EC Bext	1.7	1.2	0.3	2.2	1.2	1.8	2.2	1.0	1.9	1.3	1.1	0.9	0.8
Soil Bext	2.7	2.4	1.2	2.8	2.5	2.4	3.8	1.9	1.4	3.6	2.4	2.6	2.0
Coarse Mass Bext	8.6	6.8	4.3	8.8	8.8	9.9	10.4	5.3	4.9	11.4	5.8	6.4	5.3
Sea Salt Bext	0.1	0.3	0.2	0.0	0.1	0.3	0.0	0.2	0.2	0.4	0.2	0.3	0.2




Chiricahua - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.9	4.4	1.8	5.1	4.9	5.6	4.7	4.2	4.8	5.2	4.2	3.6	4.1
Sulfate Bext	2.3	2.1	0.3	2.5	2.2	2.5	2.2	2.1	2.4	2.3	2.0	1.7	2.3
Nitrate Bext	0.5	0.4	0.2	0.5	0.6	0.6	0.5	0.6	0.5	0.4	0.4	0.3	0.4
POM Bext	1.4	0.9	0.8	1.6	1.5	1.6	1.3	1.0	1.1	1.1	1.0	0.7	0.8
EC Bext	0.7	0.6	0.1	0.7	0.7	0.9	0.6	0.5	0.7	0.8	0.6	0.4	0.4
Soil Bext	0.3	0.3	0.1	0.4	0.2	0.5	0.3	0.2	0.3	0.5	0.3	0.3	0.4
Coarse Mass Bext	1.2	1.2	0.5	1.2	1.2	1.6	1.3	0.9	1.2	1.8	1.0	1.0	1.0
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.0




Hance Camp at Grand Canyon - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.7	12.0	7.0	11.1		11.6	12.7	11.2	12.6	11.1	11.7	10.6	13.8
Sulfate Bext	5.4	5.8	0.7	6.3		5.1	4.7	5.4	7.4	5.6	5.7	6.4	4.1
Nitrate Bext	2.2	1.8	0.7	2.4		2.1	2.0	2.3	1.3	2.1	2.0	1.6	1.8
POM Bext	10.7	10.7	5.0	5.9		9.3	17.1	10.4	13.3	7.7	8.4	4.9	19.5
EC Bext	2.4	2.9	0.4	1.6		2.1	3.1	2.8	3.7	2.4	2.1	1.3	5.3
Soil Bext	1.3	1.5	1.1	1.4		1.7	1.3	1.0	1.0	1.1	2.3	2.1	0.9
Coarse Mass Bext	3.5	3.2	3.9	3.7		3.8	3.7	2.8	2.5	2.8	3.5	3.9	3.3
Sea Salt Bext	0.1	0.1	0.2	0.4		0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.2




Hance Camp at Grand Canyon - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.2	2.2	0.3	2.9		2.0	1.8	2.0	2.1	2.4	2.5	1.6	2.2
Sulfate Bext	1.5	1.6	0.2	1.7		1.5	1.2	1.4	1.6	1.8	1.6	1.2	1.6
Nitrate Bext	0.4	0.4	0.2	0.5		0.5	0.4	0.4	0.4	0.3	0.5	0.3	0.6
POM Bext	0.6	0.5	0.3	1.0		0.4	0.5	0.6	0.5	0.6	0.6	0.4	0.4
EC Bext	0.3	0.3	0.1	0.4		0.3	0.4	0.3	0.3	0.4	0.4	0.2	0.2
Soil Bext	0.1	0.2	0.1	0.2		0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.2
Coarse Mass Bext	0.4	0.5	0.4	0.6		0.4	0.4	0.4	0.4	0.5	0.6	0.4	0.4
Sea Salt Bext	0.0	0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0




Mount Baldy - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.8	11.8	6.2	11.4	12.9	12.6	10.4	12.4	9.7	11.5	14.1	11.3	
Sulfate Bext	6.2	6.5	0.6	6.8	6.5	5.9	5.5	9.2	6.0	6.8	5.1	5.3	
Nitrate Bext	1.1	1.0	0.6	0.9	1.2	1.4	1.0	0.7	0.9	1.3	0.8	1.2	
POM Bext	13.0	10.9	4.3	9.1	22.2	12.1	8.5	11.9	4.5	8.6	17.0	12.3	
EC Bext	2.8	2.1	0.4	2.4	4.3	2.6	1.9	2.8	1.3	1.7	2.5	2.2	
Soil Bext	1.1	1.5	1.1	1.1	1.1	1.6	0.8	0.8	1.3	1.3	2.4	1.8	
Coarse Mass Bext	2.8	4.1	3.6	2.7	2.8	3.8	2.1	2.4	3.3	4.0	6.2	4.5	
Sea Salt Bext	0.0	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.2	0.1	0.1	




Mount Baldy - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.0	2.9	0.5	3.6	2.8	2.6	2.8	3.0	3.2	2.7	2.7	3.0	
Sulfate Bext	1.6	1.6	0.3	2.1	1.4	1.4	1.7	1.7	1.7	1.4	1.2	1.9	
Nitrate Bext	0.4	0.3	0.2	0.6	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	
POM Bext	1.2	1.2	0.5	1.3	1.1	1.2	1.2	1.2	1.2	1.3	1.0	1.3	1.0
EC Bext	0.6	0.6	0.1	0.6	0.7	0.6	0.5	0.7	0.7	0.7	0.6	0.5	0.5
Soil Bext	0.2	0.2	0.1	0.2	0.3	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.3
Coarse Mass Bext	0.5	0.6	0.3	0.6	0.6	0.4	0.3	0.6	0.6	0.7	0.6	0.6	0.6
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Ike's Backbone - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.3	13.4	6.7	12.1	13.5	14.5	13.3	14.7	13.0	14.3	13.2	11.9	
Sulfate Bext	6.5	7.5	0.6	6.3	7.4	6.2	6.1	9.1	6.9	8.1	7.6	5.7	
Nitrate Bext	3.5	2.3	0.7	2.2	3.1	3.6	5.2	1.4	3.4	2.7	2.2	2.0	
POM Bext	7.6	8.3	3.2	5.6	7.0	9.0	8.9	14.4	7.6	7.0	7.0	5.7	
EC Bext	2.3	2.3	0.3	2.0	2.0	2.4	3.0	3.5	2.3	1.9	1.9	1.8	
Soil Bext	2.6	2.3	1.1	2.0	3.5	3.7	1.4	1.2	1.7	3.5	2.9	2.4	
Coarse Mass Bext	6.2	6.2	3.8	5.5	6.0	9.0	4.3	5.5	4.9	9.4	5.9	5.4	
Sea Salt Bext	0.1	0.2	0.1	0.0	0.0	0.0	0.3	0.2	0.1	0.2	0.2	0.3	




Ike's Backbone - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.4	5.1	1.9	5.9	5.5	4.9	5.3	5.0	6.5	5.2	4.3	4.4	
Sulfate Bext	2.2	2.0	0.3	2.5	2.2	1.8	2.4	2.1	2.5	1.9	1.5	1.8	
Nitrate Bext	0.9	0.7	0.2	1.1	0.9	0.8	0.6	0.8	1.0	0.8	0.6	0.4	
POM Bext	1.8	1.9	0.9	1.9	1.8	1.8	1.8	1.9	2.8	1.7	1.5	1.5	
EC Bext	0.9	0.8	0.1	1.0	0.9	0.9	0.8	0.8	1.2	0.8	0.6	0.7	
Soil Bext	0.4	0.3	0.1	0.3	0.5	0.4	0.4	0.4	0.2	0.4	0.4	0.3	0.3
Coarse Mass Bext	1.0	1.0	0.5	1.2	1.2	0.8	1.0	0.7	1.3	1.3	0.8	0.8	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.0	




Petrified Forest - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.2	13.0	6.5	12.7	10.8	13.5	16.1	12.9	14.0	12.1	12.7	12.3	14.0
Sulfate Bext	6.6	7.2	0.6	6.8	6.8	6.4	6.8	6.5	8.9	6.1	7.9	6.4	6.5
Nitrate Bext	1.8	1.5	0.6	1.8	1.3	2.2	1.7	2.2	1.3	1.5	1.7	1.4	1.7
POM Bext	10.9	9.5	3.9	8.8	5.0	10.0	13.2	17.4	14.3	6.4	7.0	5.2	14.3
EC Bext	2.9	3.4	0.3	2.9	2.4	2.7	3.3	3.4	5.0	2.8	3.2	2.2	3.9
Soil Bext	2.0	2.6	1.1	1.7	1.6	2.5	3.0	1.3	2.7	2.2	2.1	3.0	2.9
Coarse Mass Bext	7.3	6.3	4.3	5.0	3.8	7.6	15.5	4.6	5.7	5.7	4.8	7.1	8.4
Sea Salt Bext	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.2	0.2	0.1




Petrified Forest - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.0	4.6	1.1	5.0	5.0	5.1	5.4	4.6	4.5	5.4	4.9	4.1	4.2
Sulfate Bext	2.3	2.1	0.3	2.2	2.4	2.3	2.1	2.2	2.3	2.4	2.3	1.6	2.1
Nitrate Bext	0.8	0.6	0.3	0.7	1.0	0.9	0.7	0.8	0.6	0.7	0.6	0.6	0.6
POM Bext	1.9	1.5	0.8	2.2	1.5	1.7	2.6	1.5	1.5	1.8	1.5	1.5	1.1
EC Bext	1.3	1.3	0.1	1.4	1.1	1.4	1.5	1.2	1.3	1.7	1.4	1.1	0.9
Soil Bext	0.4	0.4	0.2	0.3	0.5	0.5	0.3	0.3	0.4	0.6	0.4	0.3	0.4
Coarse Mass Bext	1.0	1.0	0.5	0.8	1.1	1.0	1.0	0.8	0.6	1.0	1.2	0.8	1.2
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0




Saguaro - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.8	13.6	6.5			14.8	17.2	12.5	14.3	13.9	14.1	13.2	12.7
Sulfate Bext	7.4	7.4	0.5			8.3	7.8	6.1	10.6	6.6	7.9	6.1	5.7
Nitrate Bext	5.8	2.6	0.9			6.4	6.7	4.2	2.0	3.1	3.0	2.7	2.2
POM Bext	9.4	5.3	3.2			7.3	16.0	4.8	7.2	5.8	4.7	5.4	3.4
EC Bext	3.1	2.2	0.3			2.8	3.9	2.7	3.3	2.9	1.8	1.8	1.5
Soil Bext	3.4	3.3	0.9			4.4	3.7	2.0	2.2	2.7	4.8	3.4	3.4
Coarse Mass Bext	7.1	8.3	3.3			6.2	10.1	4.9	6.8	8.9	8.4	7.8	9.6
Sea Salt Bext	0.2	0.4	0.3			0.1	0.1	0.3	0.2	0.4	0.6	0.6	0.3




Saguaro - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.9	6.7	2.2			7.4	7.1	6.4	7.2	7.8	6.6	5.9	6.2
Sulfate Bext	2.7	2.8	0.3			3.0	2.3	2.7	3.5	3.4	2.3	2.0	2.6
Nitrate Bext	1.0	0.8	0.2			0.8	1.2	0.9	1.0	0.8	0.8	0.7	0.6
POM Bext	2.2	2.0	0.9			2.2	2.3	2.0	2.4	2.2	2.1	1.8	1.5
EC Bext	1.3	1.2	0.1			1.3	1.3	1.2	1.4	1.4	1.3	0.8	0.8
Soil Bext	1.1	0.8	0.2			1.7	1.0	0.6	0.6	0.9	1.1	0.8	0.8
Coarse Mass Bext	1.8	2.1	0.7			2.0	2.0	1.5	1.8	2.9	1.8	1.7	2.3
Sea Salt Bext	0.1	0.2	0.1			0.0	0.1	0.2	0.2	0.2	0.2	0.2	0.1




Sycamore Canyon - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.3	15.2	6.7	15.0	15.5	16.3	14.2	16.7	14.3	15.8	14.7	14.4	
Sulfate Bext	5.0	5.7	0.6	4.9	4.9	4.9	5.2	8.2	5.0	4.9	6.3	4.0	
Nitrate Bext	2.0	1.4	0.6	1.1	1.8	2.2	3.0	1.2	1.0	1.6	1.1	1.9	
POM Bext	11.7	11.2	3.1	10.7	11.1	16.7	8.2	12.6	8.9	10.1	9.0	15.5	
EC Bext	3.2	3.5	0.2	2.3	3.0	4.2	3.2	3.3	3.4	2.8	3.3	4.8	
Soil Bext	6.8	5.8	1.2	7.1	9.3	5.5	5.2	6.2	5.2	8.1	5.3	4.1	
Coarse Mass Bext	9.4	10.8	4.8	11.2	8.7	9.7	7.9	14.5	9.9	12.8	9.7	6.9	
Sea Salt Bext	0.1	0.1	0.2	0.2	0.1	0.0	0.1	0.1	0.1	0.2	0.1	0.1	




Sycamore Canyon - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.6	5.1	1.0	5.5	6.1	5.9	4.9	4.8	6.3	4.7	4.5	5.4	
Sulfate Bext	2.0	2.1	0.3	2.4	2.0	1.8	2.0	2.6	2.4	1.7	1.6	2.1	
Nitrate Bext	0.9	0.8	0.3	1.2	0.9	0.7	0.7	1.0	0.8	0.8	0.6	0.9	
POM Bext	2.4	1.8	0.8	2.4	2.4	2.8	2.1	1.5	2.2	1.6	1.7	2.1	
EC Bext	1.6	1.4	0.1	1.4	1.5	1.8	1.5	0.9	1.9	1.2	1.4	1.4	
Soil Bext	0.7	0.6	0.1	0.4	1.0	1.0	0.3	0.5	0.8	0.5	0.5	0.6	
Coarse Mass Bext	1.0	1.2	0.4	0.6	1.7	1.1	0.7	0.8	1.7	1.2	0.9	1.2	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.0	




Tonto - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.2	13.8	6.5	13.9	14.8	15.2	12.8	14.3	13.9	14.7	13.7	12.6	
Sulfate Bext	7.2	8.5	0.5	7.0	7.5	6.9	7.3	11.0	8.1	9.0	7.9	6.5	
Nitrate Bext	3.0	2.5	0.7	2.9	3.2	3.1	3.0	1.8	3.2	2.7	2.7	2.1	
POM Bext	10.0	6.4	3.1	9.5	13.9	10.3	6.2	9.4	6.1	5.6	6.2	5.0	
EC Bext	2.7	2.1	0.3	3.1	3.2	2.8	1.8	3.4	2.1	1.6	1.8	1.6	
Soil Bext	2.4	2.8	1.1	1.8	2.6	3.7	1.5	1.4	2.8	3.9	3.2	2.5	
Coarse Mass Bext	7.4	7.9	3.7	7.4	6.6	9.7	6.0	5.3	8.3	11.0	7.5	7.3	
Sea Salt Bext	0.1	0.3	0.1	0.0	0.1	0.0	0.3	0.2	0.3	0.4	0.4	0.4	




Tonto - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.5	5.7	2.0	6.8	6.7	6.5	5.8	5.8	6.1	5.7	5.6	5.3	
Sulfate Bext	2.8	2.6	0.3	3.1	3.1	2.3	2.6	2.8	3.0	2.3	2.5	2.3	
Nitrate Bext	0.9	0.7	0.2	1.2	0.9	1.1	0.6	0.7	0.5	0.8	1.0	0.5	
POM Bext	2.3	1.8	0.9	2.5	2.2	2.4	2.2	2.1	2.1	1.7	1.8	1.5	
EC Bext	1.1	0.9	0.1	1.1	1.2	1.1	0.9	1.0	1.1	0.9	0.8	0.8	
Soil Bext	0.5	0.5	0.2	0.4	0.6	0.6	0.5	0.3	0.4	0.5	0.6	0.5	
Coarse Mass Bext	1.5	1.3	0.5	1.4	1.6	1.7	1.2	1.1	1.5	1.4	0.9	1.4	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.1	




Caney Creek - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	26.4	25.3	11.6			27.2	26.5	25.3	29.2	25.7		23.7	22.7
Sulfate Bext	87.1	87.0	3.2			109.5	81.7	70.0	145.8	84.7		67.8	49.7
Nitrate Bext	13.8	9.5	1.1			10.2	15.1	16.0	6.2	7.2		7.2	17.3
POM Bext	23.4	16.4	12.0			19.6	27.4	23.3	20.1	20.0		12.8	12.6
EC Bext	4.8	4.2	0.3			4.4	5.8	4.3	5.8	4.8		3.2	3.1
Soil Bext	1.1	1.2	1.1			1.4	1.1	0.9	0.7	1.6		1.9	0.8
Coarse Mass Bext	3.7	4.0	3.1			4.3	4.1	2.8	4.1	4.4		4.1	3.5
Sea Salt Bext	0.2	0.3	0.3			0.1	0.0	0.5	0.2	0.6		0.3	0.3




Caney Creek - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.2	10.7	4.2			11.9	10.7	11.1	12.9	12.5		9.2	8.1
Sulfate Bext	8.5	8.9	0.6			8.3	8.1	9.2	11.7	11.8		7.0	5.1
Nitrate Bext	4.0	3.5	0.5			5.9	2.9	3.3	5.1	4.6		2.3	2.2
POM Bext	4.3	3.2	2.0			4.5	4.2	4.3	4.5	3.6		2.6	2.2
EC Bext	1.5	1.4	0.1			1.7	1.4	1.4	2.0	1.7		1.0	0.9
Soil Bext	0.2	0.3	0.1			0.3	0.2	0.2	0.3	0.3		0.2	0.2
Coarse Mass Bext	1.8	1.9	0.9			2.0	1.8	1.6	2.4	2.4		1.3	1.4
Sea Salt Bext	0.2	0.2	0.1			0.1	0.0	0.4	0.2	0.2		0.3	0.2




Upper Buffalo - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	26.3	25.9	11.6	26.2	25.6	26.7	27.2	25.6	30.5	25.4	26.2	24.6	22.6
Sulfate Bext	83.2	90.0	3.4	89.2	73.1	100.4	84.8	68.3	169.8	75.9	85.8	69.5	48.8
Nitrate Bext	13.3	11.9	0.7	10.6	13.8	9.7	13.3	19.1	7.7	11.0	8.7	16.0	16.2
POM Bext	22.5	19.8	12.5	21.5	21.0	16.5	32.8	20.9	24.3	18.1	28.8	15.1	12.9
EC Bext	4.7	4.7	0.3	4.9	4.2	4.2	5.8	4.5	6.8	4.2	5.4	3.6	3.4
Soil Bext	1.2	1.2	1.1	1.9	1.2	1.2	0.9	0.9	0.9	1.5	1.0	1.6	0.9
Coarse Mass Bext	6.8	4.9	3.4	3.4	6.7	8.9	8.0	7.2	2.7	9.1	5.3	3.6	3.8
Sea Salt Bext	0.2	0.4	0.3	0.1	0.0	0.0	0.0	0.7	0.3	0.6	0.6	0.3	0.2




Upper Buffalo - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.7	11.7	4.2	12.9	11.5	12.8	10.6	10.7	13.3	13.0	12.4	10.5	9.4
Sulfate Bext	8.7	10.0	0.6	10.6	8.2	10.3	6.9	7.4	11.9	12.0	10.8	8.2	7.3
Nitrate Bext	4.6	3.9	0.5	6.5	4.3	5.2	2.5	4.2	6.0	4.9	3.9	3.0	1.8
POM Bext	4.2	4.0	2.0	4.8	4.5	4.6	4.1	3.2	4.7	4.4	4.7	3.3	2.9
EC Bext	1.5	1.5	0.1	1.8	1.5	1.7	1.2	1.3	2.1	1.8	1.7	1.0	1.0
Soil Bext	0.3	0.3	0.2	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.3	0.3
Coarse Mass Bext	2.8	2.3	0.8	2.3	2.5	3.4	3.5	2.4	2.4	2.5	2.7	1.9	1.9
Sea Salt Bext	0.1	0.2	0.1	0.0	0.0	0.0	0.1	0.2	0.3	0.1	0.3	0.2	0.2




Agua Tibia - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	23.5	20.9	7.6	22.9	24.9	24.0	22.2	21.2		22.0	21.0	19.4	
Sulfate Bext	31.8	25.8	1.0	29.7	34.2	36.9	26.5	33.8		24.3	23.3	22.0	
Nitrate Bext	29.9	19.9	0.9	29.5	37.6	23.5	29.1	15.2		30.8	19.1	14.3	
POM Bext	17.5	10.8	3.0	12.8	19.6	27.1	10.7	9.9		10.1	13.5	9.8	
EC Bext	6.4	4.6	0.3	6.5	6.6	7.9	4.5	6.2		4.9	3.5	3.8	
Soil Bext	1.3	0.9	0.8	1.3	1.7	1.0	1.0	0.8		0.9	0.9	1.0	
Coarse Mass Bext	8.6	7.5	3.0	8.6	10.4	7.2	8.3	6.7		8.7	8.0	6.8	
Sea Salt Bext	0.8	1.8	1.7	0.4	0.8	0.0	2.0	0.8		1.9	3.1	1.4	




Agua Tibia - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	9.6	7.4	2.9	10.2	10.4	9.4	8.4	6.9		9.0	7.5	6.3	
Sulfate Bext	3.9	2.5	0.2	3.9	4.8	3.3	3.6	2.6		3.1	2.1	2.2	
Nitrate Bext	3.3	1.6	0.2	4.5	3.8	3.1	1.7	1.2		2.2	1.6	1.3	
POM Bext	3.0	1.8	0.7	3.3	3.0	3.2	2.5	1.7		2.4	1.9	1.2	
EC Bext	1.9	1.2	0.1	2.0	2.1	2.1	1.3	1.3		1.6	1.3	0.8	
Soil Bext	0.5	0.4	0.3	0.3	0.6	0.5	0.4	0.2		0.6	0.4	0.3	
Coarse Mass Bext	2.4	2.4	0.7	2.4	2.8	2.1	2.3	1.8		3.7	2.3	1.9	
Sea Salt Bext	0.8	0.6	0.2	0.8	0.7	0.9	0.7	0.4		0.5	0.9	0.4	




Bliss SP - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.6	13.6	6.1	11.6	13.4	13.4	12.2		12.0	12.5	14.8	16.8	12.0
Sulfate Bext	5.1	6.0	0.9	5.3	4.9	4.6	5.5		6.9	6.3	5.8	6.3	4.9
Nitrate Bext	2.4	2.1	0.9	3.6	2.7	2.2	1.1		1.8	1.4	3.7	2.1	1.7
POM Bext	14.1	22.3	4.1	9.5	15.8	17.4	13.6		11.3	14.4	22.7	49.2	13.5
EC Bext	3.0	3.7	0.4	2.8	3.5	3.3	2.4		3.1	2.8	3.9	6.3	2.4
Soil Bext	1.0	0.8	1.0	0.5	1.9	1.0	0.6		0.5	0.9	0.8	0.9	0.7
Coarse Mass Bext	2.1	2.1	2.3	1.5	3.2	1.9	1.8		1.9	2.4	2.2	2.3	2.0
Sea Salt Bext	0.0	0.1	0.1	0.0	0.0	0.0	0.1		0.0	0.1	0.1	0.1	0.1




Bliss SP - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.5	2.2	0.4	3.5	2.3	2.2	2.1		2.1	2.0	2.7	2.2	1.9
Sulfate Bext	1.3	1.3	0.2	1.6	1.2	1.2	1.1		1.4	1.4	1.5	1.2	1.1
Nitrate Bext	0.5	0.4	0.2	0.7	0.6	0.4	0.4		0.4	0.3	0.5	0.3	0.3
POM Bext	1.1	0.9	0.4	1.7	0.8	0.9	1.1		0.9	0.7	1.1	1.0	0.8
EC Bext	0.5	0.3	0.1	0.6	0.5	0.4	0.4		0.3	0.3	0.3	0.2	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1		0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.4	0.4	0.3	0.4	0.3	0.4	0.3		0.3	0.4	0.5	0.5	0.4
Sea Salt Bext	0.1	0.1	0.0	0.2	0.2	0.1	0.0		0.1	0.1	0.1	0.1	0.1




Dome Land - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	19.4	19.2	7.5	18.4	21.0	19.3	19.1			20.9	19.3	17.6	
Sulfate Bext	12.0	12.8	0.9	11.7	9.8	12.8	13.8			15.0	11.4	12.0	
Nitrate Bext	15.9	15.8	0.9	17.2	20.9	13.5	12.1			19.6	13.9	14.1	
POM Bext	17.1	20.7	3.7	12.3	24.0	18.5	13.7			31.4	20.5	10.2	
EC Bext	3.9	3.7	0.4	3.5	4.7	4.1	3.4			5.4	3.3	2.4	
Soil Bext	1.5	1.7	1.0	1.7	1.9	1.1	1.3			1.9	1.7	1.4	
Coarse Mass Bext	11.0	11.7	4.2	7.5	13.1	9.8	13.6			13.3	12.0	9.8	
Sea Salt Bext	0.2	0.2	0.2	0.4	0.0	0.1	0.2			0.1	0.2	0.4	




Dome Land - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.1	5.1	1.2	5.6	4.5	5.7	4.4				5.7	4.9	4.8
Sulfate Bext	1.8	1.6	0.2	2.1	1.6	1.9	1.5				2.0	1.5	1.5
Nitrate Bext	0.9	0.9	0.1	1.2	0.7	1.3	0.6				1.2	1.0	0.7
POM Bext	2.0	1.8	0.5	2.4	1.5	2.3	1.8				1.8	2.0	1.7
EC Bext	0.8	0.7	0.1	1.0	0.9	0.9	0.6				0.7	0.7	0.8
Soil Bext	0.1	0.2	0.1	0.1	0.2	0.2	0.1				0.3	0.1	0.2
Coarse Mass Bext	0.9	1.2	0.3	0.7	0.8	1.3	0.9				1.6	0.8	1.3
Sea Salt Bext	0.1	0.2	0.1	0.1	0.1	0.0	0.1				0.2	0.3	0.1




Hoover - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.9	12.2	7.7			14.1	11.7	12.9	10.4	10.5	12.6	15.5	12.0
Sulfate Bext	5.0	5.4	0.9			4.2	4.5	6.3	5.7	5.1	5.2	5.8	4.9
Nitrate Bext	1.6	1.7	0.8			2.2	1.2	1.5	1.7	1.1	2.3	1.7	1.7
POM Bext	15.4	16.7	5.8			17.9	10.4	17.8	7.1	7.9	16.6	36.6	15.2
EC Bext	2.4	3.2	0.5			2.7	2.2	2.4	1.8	2.3	4.2	5.4	2.3
Soil Bext	1.6	1.0	1.1			2.4	1.0	1.3	0.8	1.0	1.1	1.4	1.0
Coarse Mass Bext	4.7	2.4	4.7			5.6	4.7	4.0	2.3	2.7	2.1	2.8	2.2
Sea Salt Bext	0.1	0.0	0.1			0.3	0.0	0.1	0.0	0.0	0.0	0.0	0.0




Hoover - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.4	1.3	0.1			1.8	1.1	1.4	1.1	1.3	1.7	1.2	1.1
Sulfate Bext	0.9	1.1	0.2			1.1	0.8	0.9	0.9	1.2	1.1	1.0	1.0
Nitrate Bext	0.3	0.2	0.2			0.3	0.2	0.4	0.2	0.2	0.2	0.2	0.2
POM Bext	0.5	0.5	0.2			0.4	0.5	0.5	0.5	0.3	0.8	0.5	0.3
EC Bext	0.2	0.2	0.1			0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.1
Soil Bext	0.1	0.1	0.1			0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
Coarse Mass Bext	0.5	0.3	0.3			0.7	0.3	0.4	0.3	0.4	0.4	0.2	0.4
Sea Salt Bext	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Joshua Tree - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	19.6	17.8	7.2	19.1	20.5	19.8	19.1	19.4	18.1	18.1	16.7	16.8	
Sulfate Bext	12.3	12.5	0.9	12.9	10.4	12.6	13.3	17.8	12.2	11.3	11.3	9.8	
Nitrate Bext	27.3	16.0	1.0	24.8	33.1	25.6	25.5	22.8	15.0	16.5	12.8	13.1	
POM Bext	10.3	10.3	3.5	7.6	13.6	12.0	7.8	7.7	11.1	13.8	8.4	10.5	
EC Bext	4.1	3.6	0.4	3.5	4.6	4.6	3.8	4.4	4.3	3.6	2.7	3.1	
Soil Bext	1.6	1.8	1.0	2.4	1.5	1.2	1.2	1.2	1.9	1.8	2.1	1.9	
Coarse Mass Bext	7.8	6.8	3.7	9.2	7.9	7.1	7.1	6.2	7.9	7.2	6.4	6.1	
Sea Salt Bext	0.2	0.4	0.2	0.0	0.2	0.1	0.5	0.4	0.2	0.5	0.4	0.4	




Joshua Tree - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.1	5.3	1.7	5.9	6.6	6.2	5.6	5.5	5.7	5.7	5.1	4.4	
Sulfate Bext	2.4	2.1	0.2	2.4	3.1	2.1	2.0	2.4	2.3	2.4	1.8	1.6	
Nitrate Bext	1.5	0.9	0.1	1.4	1.6	1.9	1.1	1.1	0.8	0.8	1.3	0.6	
POM Bext	1.6	1.3	0.6	1.4	1.6	1.8	1.8	1.3	1.5	1.4	1.2	1.0	
EC Bext	0.9	0.7	0.1	0.9	0.8	1.0	0.9	0.8	0.8	0.7	0.6	0.6	
Soil Bext	0.4	0.4	0.2	0.4	0.5	0.3	0.4	0.3	0.4	0.5	0.3	0.4	
Coarse Mass Bext	1.6	1.5	0.6	1.6	1.9	1.5	1.4	1.4	1.7	1.9	1.4	1.3	
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.1	0.2	0.1	0.2	0.1	




Kaiser - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.5	15.7	7.1			17.1	14.2	15.3	15.3		16.2	17.4	14.1
Sulfate Bext	7.6	8.8	1.0			7.0	6.1	9.8	11.3		8.9	7.8	7.1
Nitrate Bext	7.0	6.7	1.0			7.9	6.1	6.9	6.8		9.4	4.3	6.2
POM Bext	16.8	19.2	4.3			22.8	13.7	14.0	11.4		17.9	35.1	12.2
EC Bext	3.4	3.2	0.4			5.2	2.9	2.3	2.9		3.5	4.3	2.2
Soil Bext	1.1	1.1	0.9			1.4	0.7	1.3	0.8		1.3	1.3	1.1
Coarse Mass Bext	4.3	4.3	3.8			4.9	3.1	4.9	4.4		4.5	4.4	3.9
Sea Salt Bext	0.1	0.1	0.2			0.0	0.2	0.2	0.1		0.1	0.1	0.2




Kaiser - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.3	1.6	0.0			2.6	2.1	2.3	1.2		1.8	1.4	1.8
Sulfate Bext	1.0	1.0	0.2			1.1	1.0	1.0	0.8		1.0	1.0	1.1
Nitrate Bext	0.3	0.2	0.1			0.3	0.2	0.4	0.2		0.3	0.2	0.2
POM Bext	1.0	0.5	0.3			1.1	1.1	0.8	0.5		0.7	0.5	0.5
EC Bext	0.7	0.4	0.1			0.7	0.6	0.8	0.4		0.4	0.2	0.5
Soil Bext	0.1	0.1	0.1			0.1	0.1	0.1	0.1		0.1	0.1	0.1
Coarse Mass Bext	0.4	0.4	0.3			0.5	0.3	0.3	0.2		0.4	0.4	0.7
Sea Salt Bext	0.1	0.1	0.0			0.1	0.0	0.1	0.1		0.0	0.1	0.0




Lassen Volcanic - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.1	16.0	7.3	12.5	13.8	16.8	14.0	13.7	12.4	14.1	14.1	22.4	17.1
Sulfate Bext	6.8	7.2	1.2	6.3	7.3	6.1	6.6	7.5	8.4	7.1	7.9	6.6	5.9
Nitrate Bext	3.7	2.1	1.6	3.1	4.5	3.5	3.6	3.6	2.0	2.1	3.4	2.0	0.9
POM Bext	17.2	34.5	4.5	10.0	13.4	33.3	15.9	13.3	7.9	15.9	17.8	86.9	44.2
EC Bext	3.1	5.2	0.4	3.2	2.9	3.6	3.2	2.6	2.5	3.5	3.3	12.1	4.5
Soil Bext	1.0	0.9	0.9	0.6	1.4	1.3	0.7	0.9	0.8	1.0	0.9	1.4	0.5
Coarse Mass Bext	2.0	3.2	2.4	2.0	2.1	2.9	1.3	1.8	3.2	3.6	2.4	4.4	2.3
Sea Salt Bext	0.0	0.1	0.1	0.0	0.2	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.1




Lassen Volcanic - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.7	2.5	1.0	3.4	2.5	2.7	2.1	2.7	2.5	2.2	2.9	2.8	2.4
Sulfate Bext	1.2	1.3	0.2	1.5	1.1	1.0	0.8	1.3	1.3	1.1	1.4	1.4	1.1
Nitrate Bext	0.4	0.3	0.2	0.6	0.4	0.4	0.2	0.3	0.4	0.2	0.4	0.3	0.2
POM Bext	0.9	0.7	0.3	1.2	0.7	0.9	0.8	0.8	0.6	0.6	0.7	0.8	0.6
EC Bext	0.4	0.2	0.1	0.4	0.4	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.3	0.1	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.3	0.4	0.3
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.2	0.1	0.2	0.1	0.0




Lava Beds - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.1	14.2	7.9	14.3	15.4	16.0	14.5	13.5	14.4	12.2	16.9	13.9	
Sulfate Bext	6.8	7.6	1.4	7.6	6.5	6.5	6.6	8.2	7.4	8.5	7.1	6.8	
Nitrate Bext	3.5	2.2	1.9	3.0	5.3	3.1	2.4	3.2	1.7	2.3	2.2	1.7	
POM Bext	22.0	18.6	4.9	15.0	22.5	30.8	19.8	12.3	17.8	8.4	31.8	22.7	
EC Bext	3.3	3.1	0.5	2.8	3.3	4.2	3.1	3.2	3.2	2.2	3.9	3.1	
Soil Bext	1.0	0.8	1.0	1.6	0.9	0.7	0.8	0.5	0.9	0.8	0.7	1.0	
Coarse Mass Bext	2.6	2.3	2.8	2.9	2.3	2.5	2.9	1.9	2.7	1.9	2.5	2.4	
Sea Salt Bext	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.2	0.0	0.2	0.0	0.2	




Lava Beds - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.2	2.8	1.3	3.6	3.1	3.0	3.1	2.9	2.4	3.6	2.6	2.5	
Sulfate Bext	1.2	1.3	0.3	1.6	1.1	0.8	1.2	1.5	1.2	1.7	1.2	1.1	
Nitrate Bext	0.6	0.4	0.3	0.6	0.7	0.4	0.5	0.4	0.3	0.5	0.4	0.4	
POM Bext	1.2	0.8	0.4	1.2	1.1	1.4	1.1	0.9	0.5	1.2	0.8	0.7	
EC Bext	0.4	0.3	0.1	0.3	0.4	0.5	0.3	0.3	0.3	0.3	0.3	0.2	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.4	0.2	0.3	0.4	0.4	0.3	0.3	0.2	0.2	0.4	0.3	0.2	
Sea Salt Bext	0.1	0.2	0.0	0.1	0.0	0.0	0.2	0.2	0.2	0.2	0.2	0.2	0.2




Pinnacles - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.5	18.4	8.0	18.7		19.9	17.1	18.1	18.1	17.9	18.3	19.8	17.8
Sulfate Bext	13.9	16.6	1.0	12.4		15.4	13.8	13.9	18.2	17.3	15.7	15.5	16.1
Nitrate Bext	17.1	13.3	1.3	18.3		19.3	10.1	20.7	13.9	10.9	16.2	13.5	12.0
POM Bext	13.2	13.4	2.8	12.1		18.1	12.5	10.2	8.9	9.0	12.6	24.8	11.8
EC Bext	4.6	3.5	0.3	6.4		4.7	3.5	3.6	4.2	3.4	3.4	3.9	2.5
Soil Bext	0.6	0.6	0.6	0.6		0.6	0.5	0.6	0.4	0.8	0.5	0.6	0.6
Coarse Mass Bext	3.8	4.9	2.7	3.8		4.3	4.0	3.0	4.2	6.0	4.7	5.1	4.4
Sea Salt Bext	0.9	2.5	2.7	0.9		1.3	0.6	0.9	1.5	2.0	2.0	3.5	3.4




Pinnacles - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	8.9	8.0	3.5	9.8		8.6	8.4	8.7	8.0	7.9	8.5	8.4	7.2
Sulfate Bext	4.4	3.6	0.4	4.9		4.2	4.5	4.1	4.4	3.3	3.5	3.5	3.5
Nitrate Bext	2.6	1.8	0.3	2.8		2.6	2.6	2.4	1.7	1.5	2.2	2.1	1.5
POM Bext	2.6	2.1	0.9	2.8		2.7	2.3	2.6	2.0	2.1	2.4	2.6	1.6
EC Bext	1.2	1.0	0.1	1.5		1.2	1.1	1.2	1.1	1.1	1.1	0.8	0.7
Soil Bext	0.2	0.2	0.2	0.2		0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.2
Coarse Mass Bext	1.7	1.7	0.9	1.9		1.7	1.4	1.7	1.7	1.3	2.1	1.7	1.5
Sea Salt Bext	0.9	1.1	0.3	1.7		0.4	0.4	1.1	0.6	1.6	1.0	1.6	0.8




Point Reyes - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	22.8	22.0	15.8	21.2		24.3		22.9	22.3	22.0	22.4	22.1	21.3
Sulfate Bext	14.1	20.9	0.8	13.1		15.2		13.9	27.6	22.2	17.1	17.9	19.7
Nitrate Bext	38.4	22.5	0.9	19.9		56.6		38.7	24.8	16.0	32.4	17.7	21.7
POM Bext	12.1	5.7	2.2	8.6		16.4		11.2	5.4	4.6	5.3	7.2	5.8
EC Bext	3.3	2.0	0.2	3.9		3.1		2.8	2.6	1.9	2.2	1.5	1.6
Soil Bext	0.4	0.3	0.4	0.4		0.5		0.3	0.2	0.3	0.2	0.4	0.3
Coarse Mass Bext	7.5	7.8	3.3	7.6		7.7		7.1	5.8	9.6	8.6	9.8	5.4
Sea Salt Bext	21.2	23.4	29.7	19.8		23.2		20.6	18.5	26.0	24.0	27.6	21.2




Point Reyes - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.5	9.1	4.8	10.5		10.5		10.5	9.8	9.6	8.9	9.5	7.8
Sulfate Bext	6.7	5.6	0.8	6.8		6.1		7.3	6.0	5.7	6.0	5.9	4.4
Nitrate Bext	2.4	1.6	0.3	2.4		2.4		2.3	1.9	1.6	1.6	1.6	1.2
POM Bext	1.5	1.0	1.0	1.3		1.5		1.7	1.5	0.7	0.6	1.5	0.6
EC Bext	0.6	0.3	0.1	0.5		0.6		0.6	0.5	0.3	0.3	0.4	0.2
Soil Bext	0.1	0.1	0.1	0.1		0.1		0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	2.3	1.6	0.6	2.6		2.5		1.8	1.6	2.1	1.6	1.5	1.1
Sea Salt Bext	3.4	3.2	1.4	3.2		3.7		3.4	3.3	4.0	2.7	3.5	2.4




Redwood - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.5	19.1	13.9	19.0	18.9	18.7	17.2	18.6	18.1	20.8	18.2	19.8	18.9
Sulfate Bext	14.9	19.5	1.7	12.2	14.6	15.7	16.4	15.6	19.3	22.1	16.8	17.0	22.2
Nitrate Bext	7.0	6.2	2.0	7.5	9.8	6.5	4.5	6.4	6.5	7.5	7.4	5.1	4.4
POM Bext	8.0	8.9	2.5	8.2	6.6	11.0	5.0	9.1	4.9	9.0	4.4	15.9	10.2
EC Bext	1.5	1.2	0.2	1.9	1.4	1.8	0.7	1.5	0.9	1.5	0.9	1.4	1.2
Soil Bext	0.3	0.2	0.3	0.2	0.4	0.4	0.2	0.3	0.2	0.3	0.2	0.2	0.3
Coarse Mass Bext	4.1	3.9	3.4	4.8	3.4	4.0	3.8	4.3	3.3	5.4	3.3	4.0	3.5
Sea Salt Bext	17.6	20.4	20.0	22.0	19.5	15.2	14.8	16.7	16.4	27.6	18.9	21.7	17.6




Redwood - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.1	5.6	3.5	7.3	6.0	5.9	5.4	5.8	5.7	5.9	6.0	5.4	4.9
Sulfate Bext	2.3	2.4	0.5	2.9	2.1	2.7	1.9	2.0	2.6	2.6	2.7	2.1	2.0
Nitrate Bext	0.7	0.5	0.3	0.9	0.7	0.7	0.5	0.6	0.6	0.5	0.7	0.5	0.4
POM Bext	1.5	1.1	0.8	1.9	1.4	1.3	1.5	1.4	1.1	0.9	1.3	1.3	0.8
EC Bext	0.5	0.3	0.1	0.8	0.5	0.4	0.3	0.5	0.4	0.2	0.3	0.3	0.2
Soil Bext	0.1	0.0	0.1	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0
Coarse Mass Bext	0.9	0.8	0.6	1.2	0.8	0.8	0.9	0.8	0.7	0.9	0.8	0.8	0.7
Sea Salt Bext	1.7	1.6	0.7	2.2	1.7	1.5	1.2	1.8	1.4	2.2	1.6	1.4	1.2




San Gabriel - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	19.9	18.0	7.0			21.8	18.4	19.6	19.4	17.4	17.4	17.9	
Sulfate Bext	12.3	15.5	1.0			10.5	10.1	16.5	20.4	15.2	12.2	14.3	
Nitrate Bext	27.7	15.5	1.2			28.9	26.1	28.0	19.7	15.3	14.0	13.2	
POM Bext	15.3	11.2	3.7			25.7	9.4	10.6	10.2	9.1	12.0	13.4	
EC Bext	4.3	4.1	0.3			5.3	3.8	3.8	5.2	4.1	3.8	3.2	
Soil Bext	0.9	1.1	0.8			0.8	0.8	1.0	0.9	0.9	1.0	1.4	
Coarse Mass Bext	8.3	5.2	4.2			15.5	4.8	4.5	4.6	3.8	5.5	7.0	
Sea Salt Bext	0.4	0.4	0.5			0.3	0.4	0.6	0.6	0.3	0.3	0.6	




San Gabriel - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)			0.4			6.4	4.4	3.7	4.3	4.4	5.4	3.7	
Sulfate Bext	1.9	2.1	0.2			2.1	1.7	1.8	2.1	2.1	2.4	1.6	
Nitrate Bext	2.1	1.5	0.2			3.2	1.7	1.3	1.5	1.3	1.8	1.5	
POM Bext	1.5	1.2	0.4			2.1	1.3	1.0	1.3	1.3	1.4	0.8	
EC Bext	0.7	0.6	0.1			1.0	0.7	0.4	0.7	0.7	0.6	0.3	
Soil Bext	0.3	0.3	0.1			0.4	0.2	0.1	0.2	0.3	0.3	0.2	
Coarse Mass Bext	1.2	1.1	0.4			1.4	1.3	0.9	0.8	0.9	1.7	1.1	
Sea Salt Bext	0.1	0.2	0.1			0.1	0.0	0.1	0.1	0.2	0.2	0.3	




San Geronimo - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	22.2	20.5	7.3	21.2	23.3	22.6	21.6	21.5	19.9	21.5	20.2	19.2	
Sulfate Bext	13.2	12.0	1.0	12.5	11.2	14.6	14.5	16.0	12.4	10.8	11.5	9.3	
Nitrate Bext	44.9	30.6	0.9	37.4	58.0	45.5	39.0	37.0	31.8	27.0	31.0	26.0	
POM Bext	14.0	16.3	3.6	12.4	14.5	17.9	11.3	12.6	9.9	29.8	13.7	15.5	
EC Bext	5.2	5.2	0.3	4.9	5.8	5.4	4.6	6.5	5.3	6.3	3.8	4.1	
Soil Bext	1.1	1.2	0.9	1.2	1.1	0.9	1.2	0.9	0.9	1.5	1.4	1.0	
Coarse Mass Bext	6.1	6.2	3.9	6.2	5.4	6.4	6.6	5.4	4.5	10.4	6.3	4.4	
Sea Salt Bext	0.2	0.6	0.2	0.1	0.1	0.1	0.6	0.6	0.4	0.5	0.8	0.6	




San Gorgonio - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.4	4.5	1.2	5.0	6.2	5.1	5.3	4.1	5.0	5.5	4.4	3.6	
Sulfate Bext	1.8	1.7	0.2	1.6	2.1	1.6	2.0	1.5	2.1	2.1	1.6	1.4	
Nitrate Bext	1.8	0.9	0.1	1.7	2.6	1.7	1.3	0.6	0.8	1.3	1.2	0.6	
POM Bext	1.6	1.1	0.4	1.4	1.7	1.5	1.6	1.2	1.1	1.3	1.0	0.7	
EC Bext	1.0	0.6	0.1	1.1	1.2	0.9	0.8	0.7	0.8	0.7	0.4	0.3	
Soil Bext	0.2	0.2	0.1	0.1	0.4	0.2	0.2	0.1	0.3	0.3	0.2	0.2	
Coarse Mass Bext	0.8	1.2	0.4	0.5	0.9	0.9	1.1	1.0	1.5	1.6	0.9	1.1	
Sea Salt Bext	0.2	0.1	0.1	0.3	0.0	0.2	0.2	0.1	0.1	0.1	0.3	0.1	




San Rafael - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.8	19.2	7.6	18.9	17.9	19.7	18.7	19.0	18.7	20.2	20.6	19.6	17.0
Sulfate Bext	20.4	20.2	1.1	22.6	20.0	18.4	19.8	21.3	22.7	25.9	17.7	18.8	15.9
Nitrate Bext	12.6	11.6	1.0	13.6	11.4	14.8	10.0	13.1	13.1	11.8	14.5	9.2	9.2
POM Bext	12.4	17.9	3.3	8.6	7.8	19.5	15.2	10.8	8.9	21.8	29.7	21.5	7.7
EC Bext	3.0	4.5	0.3	3.6	2.6	3.3	3.2	2.4	3.0	9.6	5.4	3.0	1.7
Soil Bext	1.0	1.1	1.0	0.9	1.4	0.9	1.1	1.0	0.7	1.1	0.9	1.5	1.1
Coarse Mass Bext	6.9	7.0	3.2	6.5	7.3	6.1	6.8	7.7	6.0	7.5	7.1	7.2	7.2
Sea Salt Bext	0.8	1.8	1.5	0.8	0.1	0.8	0.5	1.5	1.1	1.8	1.0	2.1	3.0




San Rafael - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.4	5.5	1.8	7.1	5.5	7.1	5.7	6.7	5.0	5.2	6.8	5.9	4.6
Sulfate Bext	2.8	2.5	0.2	3.2	2.0	3.0	2.5	3.1	2.3	2.5	3.0	2.7	2.2
Nitrate Bext	2.1	1.4	0.3	2.3	2.1	2.7	1.3	2.0	1.1	1.3	2.4	1.5	1.0
POM Bext	1.8	1.2	0.6	1.9	1.4	2.0	1.8	2.0	1.4	1.0	1.3	1.3	0.8
EC Bext	0.7	0.5	0.1	0.9	0.6	0.8	0.6	0.7	0.6	0.4	0.5	0.5	0.3
Soil Bext	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Coarse Mass Bext	1.3	1.2	0.5	1.6	1.2	1.4	1.2	1.1	0.8	0.9	1.9	1.1	1.0
Sea Salt Bext	0.5	0.7	0.1	0.7	0.3	0.6	0.3	0.9	0.6	0.7	0.8	1.1	0.5




Sequoia - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	25.4	23.4	7.7	25.7	24.5	27.0	26.0	23.7	23.1	23.4	24.7	24.7	21.0
Sulfate Bext	16.5	17.0	0.8	20.1	14.5	14.2	15.7	18.2	20.0	22.3	14.8	14.2	13.7
Nitrate Bext	60.7	45.6	1.2	76.1	63.5	66.7	43.3	54.0	40.5	44.9	66.5	34.2	42.0
POM Bext	32.4	26.4	3.3	26.8	20.6	43.9	51.6	19.2	20.6	21.3	21.2	54.3	14.7
EC Bext	7.0	5.7	0.3	7.9	5.8	7.7	8.9	4.7	5.9	5.6	6.0	7.5	3.7
Soil Bext	1.1	1.0	1.0	0.8	0.8	1.1	1.4	1.2	0.6	1.1	0.9	1.3	1.1
Coarse Mass Bext	7.7	5.8	3.5	5.6	7.0	9.4	9.2	7.1	5.5	5.1	6.1	6.4	5.8
Sea Salt Bext	0.2	0.3	0.6	0.1	0.3	0.3	0.1	0.5	0.2	0.4	0.3	0.2	0.5




Sequoia - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	8.8	7.9	2.3	9.7	7.8	8.9	8.5	8.9	7.2	7.7	9.2	9.3	6.2
Sulfate Bext	3.0	3.2	0.2	3.3	2.8	3.2	2.7	2.8	3.1	3.6	3.4	3.5	2.2
Nitrate Bext	3.5	2.7	0.2	3.7	2.5	3.9	3.5	3.7	1.6	2.5	3.9	4.3	1.2
POM Bext	3.9	2.9	0.5	5.0	2.9	3.7	4.0	3.7	2.4	2.6	3.5	4.1	2.0
EC Bext	1.5	1.0	0.1	2.0	1.1	1.3	1.5	1.4	1.1	0.9	1.3	1.1	0.7
Soil Bext	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.1	0.2	0.3	0.3	0.2
Coarse Mass Bext	1.6	1.4	0.4	1.9	1.5	1.5	1.4	1.7	1.3	1.1	1.9	1.7	1.3
Sea Salt Bext	0.2	0.3	0.1	0.2	0.3	0.1	0.0	0.2	0.2	0.3	0.3	0.4	0.1




Trinity - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.3	17.3	7.9		18.1	20.5	15.2	15.7		19.7	13.9	23.1	12.7
Sulfate Bext	8.4	9.8	1.6		8.6	7.8	9.0	8.1		9.6	9.7	10.6	9.4
Nitrate Bext	6.1	3.4	2.4		10.3	5.0	3.5	5.4		4.3	3.0	5.0	1.4
POM Bext	35.3	56.8	4.7		32.8	59.0	18.0	31.2		67.8	11.8	139.0	8.8
EC Bext	4.6	6.9	0.5		5.3	5.5	3.2	4.5		9.9	2.7	13.3	1.5
Soil Bext	0.8	0.9	0.9		1.1	0.7	0.6	1.0		1.0	0.8	1.0	0.7
Coarse Mass Bext	2.5	3.6	2.7		3.3	3.1	2.5	1.3		3.4	3.8	3.7	3.5
Sea Salt Bext	0.3	0.3	0.4		0.0	0.3	0.7	0.2		0.1	0.2	0.5	0.4




Trinity - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.4	3.2	1.2	3.4	4.0	2.9	3.4		2.1	3.9	4.0	2.6	
Sulfate Bext	1.3	1.6	0.3	1.5	1.5	0.9	1.2		0.8	2.1	2.1	1.3	
Nitrate Bext	0.5	0.4	0.2	0.6	0.6	0.3	0.4		0.3	0.5	0.4	0.3	
POM Bext	1.4	0.9	0.3	1.1	1.7	1.3	1.5		0.6	1.0	1.4	0.9	
EC Bext	0.5	0.3	0.1	0.4	0.5	0.5	0.4		0.2	0.3	0.3	0.2	
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.0	0.1		0.0	0.1	0.1	0.1	
Coarse Mass Bext	0.3	0.3	0.2	0.3	0.3	0.2	0.2		0.1	0.4	0.3	0.2	
Sea Salt Bext	0.2	0.4	0.1	0.1	0.3	0.2	0.4		0.4	0.5	0.6	0.1	




Yosemite - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.6	16.9	7.6	16.6	16.3	20.2	18.0	17.1	15.7	15.9	17.7	18.6	16.5
Sulfate Bext	7.9	9.2	0.9	9.5	7.7	7.4	6.4	8.5	10.2	10.9	8.2	8.7	8.0
Nitrate Bext	8.1	6.5	1.2	15.6	7.4	8.3	5.7	3.5	4.6	5.5	13.4	3.2	5.8
POM Bext	29.0	26.7	4.7	16.1	19.0	42.4	35.5	31.8	16.4	14.5	25.5	50.2	26.6
EC Bext	4.8	4.6	0.5	4.6	4.0	5.5	5.2	4.6	3.8	3.6	5.0	6.7	4.2
Soil Bext	0.9	0.9	0.9	0.8	1.2	0.8	0.6	1.1	0.5	1.3	0.8	1.1	0.9
Coarse Mass Bext	4.0	4.0	3.7	4.5	3.6	4.0	3.4	4.4	3.5	3.6	4.2	4.5	4.0
Sea Salt Bext	0.1	0.1	0.1	0.0	0.0	0.1	0.3	0.1	0.1	0.2	0.1	0.1	0.3




Yosemite - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.4	2.9	1.0	4.2	2.8	3.7	3.2	3.1	3.2	3.4	3.9	2.0	2.3
Sulfate Bext	1.4	1.4	0.2	1.8	1.2	1.4	1.2	1.2	1.3	1.7	1.8	1.0	1.2
Nitrate Bext	0.6	0.5	0.1	0.5	0.6	0.7	0.5	0.5	0.8	0.4	0.6	0.2	0.2
POM Bext	1.2	0.8	0.3	1.6	0.6	1.2	1.3	1.1	0.9	0.9	1.1	0.4	0.4
EC Bext	0.4	0.3	0.0	0.5	0.2	0.4	0.3	0.4	0.4	0.4	0.4	0.1	0.2
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
Coarse Mass Bext	0.5	0.4	0.3	0.5	0.5	0.5	0.4	0.4	0.2	0.3	0.7	0.3	0.5
Sea Salt Bext	0.1	0.1	0.0	0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.0




Great Sand Dunes - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.8	11.4	6.7	14.1	11.4	14.3	12.9	11.2	12.5	11.6	11.0	11.4	10.8
Sulfate Bext	6.0	6.6	0.7	6.2	7.0	5.1	6.3	5.2	8.2	7.1	6.0	6.0	5.9
Nitrate Bext	2.0	1.5	0.9	1.7	1.8	2.5	1.1	2.7	1.2	1.0	2.6	1.6	1.4
POM Bext	8.5	6.1	3.4	10.6	5.1	10.8	9.1	6.8	7.5	5.9	6.3	5.5	5.2
EC Bext	1.7	1.7	0.3	2.2	1.2	2.0	1.9	1.3	2.1	2.0	1.5	1.6	1.3
Soil Bext	2.8	1.9	1.2	2.8	2.3	4.1	2.8	2.1	1.9	1.8	0.9	2.5	2.5
Coarse Mass Bext	7.2	5.1	4.2	9.5	5.8	10.8	6.5	3.7	5.9	5.6	4.0	5.8	4.4
Sea Salt Bext	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.2	0.0	0.2	0.1	0.1	0.1




Great Sand Dunes - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.5	3.6	1.2	4.5	4.5	5.0	4.5	4.1	3.3	3.9	3.6	3.4	3.6
Sulfate Bext	2.1	1.9	0.3	2.4	2.2	2.4	1.7	1.9	1.6	2.2	1.9	1.7	2.1
Nitrate Bext	0.7	0.5	0.3	0.6	0.5	0.9	0.6	0.7	0.4	0.4	0.6	0.6	0.5
POM Bext	1.8	1.4	0.8	1.6	1.5	1.8	2.2	1.9	1.4	1.2	1.4	1.4	1.5
EC Bext	0.6	0.5	0.1	0.6	0.6	0.7	0.5	0.6	0.5	0.7	0.5	0.5	0.6
Soil Bext	0.4	0.3	0.2	0.3	0.5	0.4	0.4	0.3	0.2	0.4	0.3	0.2	0.3
Coarse Mass Bext	1.1	0.7	0.5	1.1	1.3	1.3	1.2	0.6	0.7	0.9	0.7	0.7	0.5
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Mesa Verde - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.0	11.3	6.8	13.0	10.5	14.7	16.8	10.3	12.0	10.7	10.9	11.0	11.7
Sulfate Bext	6.5	6.3	0.7	6.6	7.3	5.5	5.9	7.0	7.6	6.2	6.5	6.3	4.9
Nitrate Bext	2.3	2.0	0.8	1.7	1.9	3.5	2.5	2.0	1.5	1.6	3.1	1.3	2.4
POM Bext	12.3	6.5	4.2	17.8	4.1	11.5	23.4	4.7	8.9	6.2	6.3	5.2	5.9
EC Bext	2.4	1.6	0.4	3.6	1.3	2.2	3.5	1.2	2.6	1.7	1.6	1.0	1.3
Soil Bext	2.5	2.0	1.2	1.9	1.6	3.9	3.9	1.3	1.5	1.9	1.1	2.3	3.2
Coarse Mass Bext	6.5	4.6	4.3	3.8	3.5	9.5	12.8	3.0	4.0	3.2	2.4	5.6	7.5
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.1	0.1	0.0




Mesa Verde - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.3	3.1	1.0	4.3	4.0	4.7	4.7	3.9	3.0	3.6	3.3	2.8	3.0
Sulfate Bext	2.4	2.1	0.4	2.3	2.7	2.5	2.1	2.3	2.0	2.3	2.1	1.8	2.2
Nitrate Bext	0.8	0.6	0.4	0.8	0.7	1.0	1.0	0.8	0.6	0.6	0.6	0.6	0.6
POM Bext	1.5	1.0	0.7	1.8	1.0	1.3	1.9	1.5	1.0	1.1	1.2	1.0	0.8
EC Bext	0.6	0.4	0.1	0.8	0.4	0.6	0.7	0.6	0.4	0.6	0.4	0.3	0.3
Soil Bext	0.4	0.2	0.2	0.2	0.4	0.6	0.5	0.3	0.3	0.3	0.2	0.2	0.2
Coarse Mass Bext	0.7	0.4	0.4	0.6	0.9	1.1	0.9	0.4	0.3	0.4	0.5	0.5	0.5
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Mount Zirkel - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.5	9.7	6.1	10.3	12.5	10.6	8.7	9.4	9.7	10.1	9.7	9.5	
Sulfate Bext	5.2	5.6	0.7	6.8	4.9	4.7	4.6	7.0	5.6	5.2	5.5	4.7	
Nitrate Bext	2.2	1.4	0.9	2.9	2.3	1.5	1.9	1.3	1.0	1.4	1.9	1.5	
POM Bext	9.9	7.2	4.5	5.1	17.3	11.6	5.8	4.8	7.2	8.9	5.9	9.4	
EC Bext	1.8	1.5	0.4	1.4	2.6	2.0	1.0	1.5	1.7	1.7	1.0	1.7	
Soil Bext	1.0	1.1	1.0	1.2	1.2	0.6	1.0	0.9	0.7	1.0	1.4	1.8	
Coarse Mass Bext	3.1	2.9	3.6	3.2	3.8	3.8	1.8	2.4	2.6	2.1	3.5	3.9	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	




Mount Zirkel - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.6	0.7	-0.5	2.7	1.4	1.2	1.1	0.7	1.0	0.8	0.4	0.6	
Sulfate Bext	1.5	1.2	0.3	2.3	1.3	1.2	1.3	1.2	1.4	1.2	1.1	1.2	
Nitrate Bext	0.5	0.4	0.2	0.6	0.5	0.6	0.4	0.4	0.4	0.4	0.4	0.5	0.4
POM Bext	0.8	0.5	0.5	1.1	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.4	0.5
EC Bext	0.3	0.2	0.1	0.4	0.4	0.3	0.3	0.2	0.3	0.3	0.3	0.1	0.2
Soil Bext	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.5	0.2	0.4	0.6	0.5	0.5	0.5	0.2	0.3	0.2	0.2	0.2	0.2
Sea Salt Bext	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Rocky Mountain National Park - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.8	12.6	7.2	14.0	13.5	16.0	13.7	11.9	13.5	12.5	12.6	12.8	11.7
Sulfate Bext	7.9	7.2	0.9	6.4	9.7	9.4	6.8	7.3	9.3	7.3	6.9	5.9	6.5
Nitrate Bext	5.3	4.0	1.0	2.5	7.6	8.7	3.2	4.3	4.8	3.7	4.3	4.1	3.2
POM Bext	10.5	8.9	4.3	11.1	6.2	16.5	12.1	6.7	7.8	9.0	8.5	9.1	10.1
EC Bext	2.6	2.2	0.4	2.9	2.0	3.4	2.5	2.0	2.5	2.4	1.9	1.7	2.3
Soil Bext	1.4	1.5	1.1	1.3	1.4	1.9	1.2	1.1	1.5	0.9	1.3	2.3	1.5
Coarse Mass Bext	4.9	3.9	4.1	8.7	3.2	4.6	5.1	2.9	4.1	3.2	3.8	5.1	3.2
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1




Rocky Mountain National Park - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.3	2.0	0.3	2.6	2.4	1.8	2.5	2.1	2.1	2.1	2.3	1.6	2.0
Sulfate Bext	1.3	1.2	0.2	1.7	1.5	0.9	1.3	1.2	1.4	1.3	1.2	1.0	1.3
Nitrate Bext	0.3	0.3	0.1	0.5	0.3	0.2	0.3	0.4	0.4	0.2	0.3	0.2	0.2
POM Bext	0.9	0.8	0.4	0.8	0.7	0.8	1.1	1.0	0.7	0.8	1.0	0.7	0.7
EC Bext	0.3	0.3	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2
Soil Bext	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.3
Coarse Mass Bext	0.6	0.6	0.4	0.6	0.8	0.6	0.8	0.4	0.5	0.6	0.7	0.5	0.5
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Weminuche - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.3	10.0	6.2	9.6	11.2	11.3	9.3	10.0	9.2	10.3	10.8	10.0	
Sulfate Bext	5.0	5.1	0.7	6.2	4.5	4.9	4.4	7.0	4.4	5.1	4.9	4.2	
Nitrate Bext	1.2	1.0	0.9	1.0	1.3	1.7	0.9	0.8	0.8	1.1	1.2	1.2	
POM Bext	8.3	6.9	3.7	4.5	9.5	12.7	6.6	5.1	5.4	7.9	9.6	6.6	
EC Bext	2.0	1.8	0.3	2.0	1.9	2.2	2.0	2.1	1.8	2.0	1.8	1.5	
Soil Bext	1.3	1.3	1.2	1.2	1.5	1.4	0.8	0.9	0.9	1.1	0.9	1.5	2.1
Coarse Mass Bext	3.0	3.0	3.3	2.1	4.1	3.6	2.2	2.2	2.7	2.8	3.6	3.6	
Sea Salt Bext	0.1	0.0	0.1	0.3	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.0	




Weminuche - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.1	2.4	1.0	3.2	3.3	2.9	3.0	2.6	2.9	2.3	2.2	2.1	
Sulfate Bext	1.5	1.3	0.3	1.7	1.6	1.2	1.4	1.4	1.7	1.2	1.1	1.4	
Nitrate Bext	0.3	0.2	0.2	0.3	0.4	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.2
POM Bext	1.2	0.9	0.7	1.2	1.2	1.2	1.3	1.2	1.0	0.9	0.7	0.6	
EC Bext	0.8	0.5	0.1	0.8	0.7	0.8	0.9	0.7	0.8	0.5	0.5	0.3	
Soil Bext	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.2
Coarse Mass Bext	0.7	0.6	0.5	0.6	0.7	0.7	0.5	0.4	0.6	0.6	0.6	0.6	0.6
Sea Salt Bext	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0




White River - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	9.6	8.9	6.1	9.4	11.8	9.6	7.6	9.1	8.8	9.0	9.6	8.3	
Sulfate Bext	4.8	5.1	0.9	6.1	4.3	4.6	4.0	6.0	5.1	4.9	5.1	4.4	
Nitrate Bext	1.3	1.3	0.9	1.5	1.5	1.1	1.1	1.0	1.2	1.5	1.7	1.3	
POM Bext	7.8	5.6	4.6	4.4	15.2	7.2	4.5	5.2	5.1	6.4	6.2	5.0	
EC Bext	1.8	1.4	0.4	1.6	2.5	1.5	1.4	1.8	1.5	1.4	1.2	1.2	
Soil Bext	1.2	1.3	1.2	1.6	1.3	0.9	0.9	1.0	1.1	1.0	1.6	1.6	
Coarse Mass Bext	2.8	2.3	3.0	2.8	3.5	3.1	1.8	2.1	2.3	1.9	2.9	2.2	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	




White River - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	0.7	0.2	-0.8	1.1	0.6	0.6	0.6	-0.1	0.5	0.3	0.1	0.1	
Sulfate Bext	1.1	1.0	0.2	1.3	0.9	1.0	1.1	0.9	1.3	0.9	1.0	1.0	
Nitrate Bext	0.3	0.2	0.2	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3	0.2
POM Bext	0.6	0.3	0.4	0.7	0.7	0.5	0.5	0.3	0.4	0.4	0.4	0.3	0.1
EC Bext	0.5	0.3	0.1	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.3	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.2	0.4
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Chassahowitzka - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	25.8	23.7	11.0	27.3	25.8	25.2	24.8	25.7		26.0	25.1	22.5	21.1
Sulfate Bext	87.6	64.1	2.8	99.7	90.0	78.6	75.4	94.3		85.9	72.4	50.8	47.4
Nitrate Bext	5.2	4.9	1.2	6.4	5.6	5.3	4.7	3.7		4.4	5.0	5.6	4.7
POM Bext	20.7	17.8	11.4	28.2	19.3	20.6	17.8	17.4		19.3	27.2	13.8	11.0
EC Bext	6.7	6.4	0.3	8.7	5.8	6.2	7.0	5.9		8.9	7.3	4.8	4.7
Soil Bext	0.9	1.3	0.8	0.7	1.4	0.7	0.9	0.8		1.4	0.9	2.0	0.7
Coarse Mass Bext	3.0	3.6	2.0	3.0	3.1	2.7	3.4	2.7		4.4	3.0	4.4	2.7
Sea Salt Bext	0.2	1.5	1.1	0.2	0.4	0.0	0.1	0.4		0.6	0.8	3.1	1.7




Chassahowitzka - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.5	15.4	5.9	16.4	16.3	15.0	14.8	15.0		16.9	15.8	15.0	14.0
Sulfate Bext	18.3	17.7	1.3	20.9	21.5	17.2	15.8	16.2		21.2	17.8	16.7	15.4
Nitrate Bext	4.2	4.2	0.8	4.5	5.2	4.4	3.0	3.7		4.8	4.5	4.3	3.3
POM Bext	6.3	5.7	3.0	6.6	5.2	5.7	7.2	6.9		6.5	6.5	5.2	4.6
EC Bext	2.9	2.5	0.1	3.5	2.6	2.4	3.5	2.6		3.5	2.5	2.1	1.7
Soil Bext	0.6	0.5	0.3	0.4	0.9	0.3	0.4	0.7		0.4	0.5	0.4	0.5
Coarse Mass Bext	3.3	3.0	1.5	3.1	3.6	3.6	3.2	3.3		3.8	3.3	2.7	2.2
Sea Salt Bext	1.0	3.0	0.2	1.6	1.5	0.6	0.6	0.8		3.4	3.1	3.1	2.3




Everglades - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	22.3	20.4	12.1		20.9	20.6	24.4	23.2	22.3	21.4	21.0	19.2	18.1
Sulfate Bext	42.5	41.9	1.7		42.2	39.1	46.5	42.2	60.0	49.8	39.2	33.9	26.6
Nitrate Bext	4.8	3.8	0.9		6.4	3.8	4.7	4.4	4.1	3.6	3.5	4.0	3.7
POM Bext	47.1	10.5	21.7		12.8	18.1	113.1	44.2	11.8	9.8	20.1	6.1	4.8
EC Bext	5.9	3.4	0.3		4.9	4.7	9.6	4.4	4.5	4.1	4.7	1.9	1.6
Soil Bext	0.9	2.3	1.1		0.8	1.3	0.9	0.8	0.6	2.8	1.5	2.7	3.7
Coarse Mass Bext	3.0	5.2	2.5		3.6	3.2	2.6	2.5	3.1	6.0	5.0	5.8	5.9
Sea Salt Bext	1.2	2.5	3.8		0.7	0.7	0.2	3.2	1.4	1.1	1.6	4.3	4.1




Everglades - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.7	11.9	5.2	11.4	11.0	12.2	12.1	12.6	13.1	11.4	11.5	11.0	
Sulfate Bext	9.6	10.2	1.0	8.4	9.1	10.8	10.0	11.8	11.8	9.2	9.0	9.2	
Nitrate Bext	2.2	2.4	0.5	2.2	2.0	2.1	2.5	2.4	2.6	2.5	2.3	2.2	
POM Bext	3.5	3.0	2.3	3.0	3.0	4.3	3.8	2.6	2.8	3.2	3.6	2.7	
EC Bext	1.1	1.1	0.1	1.3	1.0	1.3	1.0	1.1	1.3	1.1	0.9	0.9	
Soil Bext	0.4	0.4	0.2	0.4	0.4	0.4	0.4	0.4	0.3	0.8	0.3	0.5	0.3
Coarse Mass Bext	2.7	2.8	1.1	3.2	2.3	2.7	2.5	2.9	4.2	2.9	2.3	1.9	
Sea Salt Bext	1.9	2.3	0.7	2.0	1.3	1.6	2.6	3.1	2.7	1.4	2.1	2.2	




Saint Marks - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	26.3	25.6	11.7		25.9	26.1	26.0	27.3		27.9	26.8	24.6	23.0
Sulfate Bext	93.1	81.5	2.8		91.7	92.1	89.4	99.3		122.0	83.2	66.7	54.3
Nitrate Bext	4.5	4.6	1.4		4.9	4.8	4.2	4.1		3.8	4.8	4.5	5.4
POM Bext	24.2	27.8	13.0		19.7	20.6	22.7	33.9		19.3	42.5	26.6	22.7
EC Bext	5.6	5.9	0.3		5.0	6.0	6.0	5.6		6.7	6.1	4.8	5.8
Soil Bext	0.8	1.0	0.8		0.7	0.9	0.8	0.7		0.7	1.6	0.8	1.0
Coarse Mass Bext	2.6	3.4	2.3		2.9	3.1	2.9	1.5		2.6	5.2	3.9	2.1
Sea Salt Bext	0.2	1.4	1.4		0.3	0.0	0.0	0.7		1.0	0.8	2.7	1.1




Saint Marks - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.3	14.8	5.4	15.0	14.1	13.7	14.4			16.2	14.7	14.8	13.4
Sulfate Bext	18.6	19.1	1.2	19.9	18.5	16.9	19.0			25.4	17.6	18.2	15.2
Nitrate Bext	2.8	3.0	0.6	3.4	2.8	2.5	2.3			3.0	3.1	3.6	2.4
POM Bext	4.9	5.0	2.4	5.5	4.4	4.7	5.1			5.4	5.4	4.6	4.6
EC Bext	1.7	1.7	0.1	2.2	1.6	1.5	1.7			2.0	1.9	1.4	1.4
Soil Bext	0.5	0.5	0.2	0.5	0.5	0.4	0.8			0.3	0.5	0.7	0.4
Coarse Mass Bext	2.3	2.0	1.3	2.3	2.5	2.4	2.0			1.9	2.3	2.1	1.7
Sea Salt Bext	0.6	2.4	0.3	0.7	0.2	0.5	1.0			2.7	2.2	2.6	2.2




Cohutta - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	30.3	28.0	10.8	30.1	30.9	30.3	29.7	31.1	31.1	30.5	27.2	23.2	
Sulfate Bext	171.6	134.4	4.0	164.2	184.6	179.7	157.9	193.8	193.8	163.1	118.8	61.7	
Nitrate Bext	3.5	5.2	0.8	3.8	4.6	3.0	2.6	5.5	5.5	4.2	2.2	9.1	
POM Bext	22.1	20.5	10.3	22.1	25.9	19.5	21.0	18.8	18.8	31.8	17.1	14.4	
EC Bext	5.1	4.7	0.3	4.9	5.0	5.4	5.0	6.2	6.2	5.3	3.4	4.0	
Soil Bext	1.0	0.8	1.0	0.9	1.1	0.8	1.0	0.7	0.7	0.9	1.0	0.6	
Coarse Mass Bext	1.8	2.5	2.3	2.2	1.7	1.7	1.5	2.1	2.1	3.3	2.1	2.5	
Sea Salt Bext	0.1	0.1	0.2	0.1	0.0	0.0	0.2	0.1	0.1	0.2	0.2	0.1	




Cohutta - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.7	12.5	4.3	14.9	13.9	12.8	13.3	14.3		13.9	12.2	9.6	
Sulfate Bext	16.1	13.1	0.7	18.4	17.1	14.0	15.1	17.3		14.7	12.2	8.3	
Nitrate Bext	5.2	4.9	0.7	6.3	4.6	5.0	4.8	5.7		7.4	4.6	2.1	
POM Bext	4.8	3.8	2.0	5.9	4.5	4.3	4.5	4.7		4.4	3.5	2.8	
EC Bext	2.0	1.6	0.1	2.4	1.8	1.9	1.9	2.2		1.7	1.5	1.0	
Soil Bext	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2		0.2	0.2	0.2	
Coarse Mass Bext	1.0	1.1	0.6	1.2	1.3	0.8	0.7	1.0		1.0	1.2	1.3	
Sea Salt Bext	0.1	0.2	0.1	0.4	0.0	0.0	0.2	0.1		0.4	0.4	0.0	




Okefenokee - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	27.1	26.0	11.4	28.5	27.4	27.1	25.5	27.1	28.6	27.9	26.9	23.9	22.7
Sulfate Bext	102.9	94.0	3.3	108.9	98.6	96.0	86.4	124.4	135.5	116.5	85.3	72.7	60.1
Nitrate Bext	4.9	4.0	1.3	5.7	5.3	5.6	4.7	3.1	2.9	3.6	5.1	3.7	4.4
POM Bext	30.3	28.8	12.0	41.4	36.2	35.1	21.1	17.7	24.5	26.2	60.7	17.8	14.6
EC Bext	6.2	5.6	0.3	8.3	5.6	7.5	5.1	4.4	6.2	6.6	7.4	3.9	4.0
Soil Bext	0.8	1.0	0.8	0.8	0.8	0.8	0.9	1.0	0.7	0.8	1.6	0.8	1.1
Coarse Mass Bext	3.1	3.1	2.5	5.0	3.1	2.7	2.4	2.1	2.9	2.9	5.1	2.0	2.9
Sea Salt Bext	0.2	0.6	1.3	0.3	0.3	0.1	0.0	0.6	0.6	0.4	0.6	1.1	0.5




Okefenokee - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.2	14.5	5.4	16.9	15.0	15.2	13.5	15.6	15.5	15.4	14.6	13.5	13.4
Sulfate Bext	19.7	18.1	1.2	24.3	17.2	19.1	16.0	21.8	20.8	21.0	17.5	15.7	15.8
Nitrate Bext	3.7	2.8	0.7	4.9	3.9	4.0	2.1	3.3	2.7	3.3	2.8	2.5	2.4
POM Bext	6.8	5.2	2.7	8.0	7.2	7.1	6.0	5.9	6.1	5.2	5.6	4.6	4.6
EC Bext	2.2	1.9	0.1	2.8	2.1	2.2	1.7	2.0	2.6	2.0	2.2	1.4	1.3
Soil Bext	0.4	0.4	0.2	0.4	0.5	0.4	0.4	0.5	0.3	0.5	0.4	0.5	0.4
Coarse Mass Bext	2.2	2.0	1.0	3.0	2.3	2.0	1.9	1.8	2.4	2.3	2.2	1.4	1.9
Sea Salt Bext	0.7	1.6	0.2	0.3	1.1	0.3	0.0	1.8	1.3	1.9	1.7	1.8	1.5




Haleakala - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.3	14.8	7.4	13.1	12.6	14.8	12.8	13.5	13.7	13.3	17.9	15.4	
Sulfate Bext	17.5	26.5	0.9	16.2	15.2	23.5	15.3	18.8	19.4	18.6	45.3	30.3	
Nitrate Bext	2.7	2.1	1.2	3.1	2.8	2.4	2.4	2.3	2.4	2.4	1.7	1.8	
POM Bext	2.9	2.2	3.0	2.2	3.0	3.6	2.8	2.3	2.0	2.0	1.7	3.0	
EC Bext	1.4	1.2	0.4	1.2	1.3	1.7	1.3	1.3	1.0	1.0	1.1	1.3	
Soil Bext	0.3	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.4	
Coarse Mass Bext	2.6	1.9	2.7	3.2	2.7	2.4	2.2	1.9	2.3	2.0	1.6	1.9	
Sea Salt Bext	1.3	2.0	2.7	1.4	0.7	1.3	1.7	2.1	2.5	2.5	1.3	1.5	




Haleakala - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.5	4.4	2.7	4.6	4.8	4.3	4.5	3.7	4.5	4.6	4.6	4.6	4.6
Sulfate Bext	2.2	2.1	0.3	2.3	2.4	2.1	2.0	1.7	2.3	2.2	2.2	2.2	2.1
Nitrate Bext	0.6	0.4	0.2	0.5	0.7	0.5	0.5	0.3	0.4	0.5	0.4	0.4	0.5
POM Bext	0.7	0.5	0.7	0.4	0.7	0.7	0.9	0.6	0.4	0.6	0.6	0.5	0.5
EC Bext	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.2
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1
Coarse Mass Bext	1.0	0.9	1.0	1.2	1.1	1.0	0.9	0.8	0.8	0.9	0.9	0.9	1.0
Sea Salt Bext	1.1	1.5	0.7	1.1	1.0	0.9	1.3	1.1	1.4	1.6	1.7	1.7	1.7




Hawaii Volcanoes - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.9	24.9	7.2		16.4	18.0	22.0	19.0	22.1	20.6	23.8	27.5	30.5
Sulfate Bext	60.0	132.1	1.7		41.7	53.0	89.3	56.2	85.2	69.2	108.6	179.5	218.1
Nitrate Bext	0.9	0.6	1.3		1.0	1.0	0.8	0.6	0.6	0.3	0.7	0.7	0.5
POM Bext	2.5	1.3	2.0		3.6	3.9	1.3	1.2	1.4	1.0	1.5	1.7	0.9
EC Bext	1.0	0.8	0.3		0.9	0.9	1.2	1.0	0.9	0.8	0.9	0.9	0.6
Soil Bext	0.3	0.5	0.3		0.2	0.2	0.4	0.3	0.3	0.2	0.3	0.7	1.2
Coarse Mass Bext	0.8	1.1	1.3		1.1	0.4	0.7	0.9	1.0	0.4	0.8	2.5	0.8
Sea Salt Bext	1.2	1.3	3.7		2.3	0.4	0.1	1.4	1.5	1.5	1.2	1.5	0.8




Hawaii Volcanoes - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.1	3.8	2.2	4.1	3.9	4.2	4.0	3.5	3.2	4.0	4.1	3.9	
Sulfate Bext	2.2	2.2	0.2	1.9	1.7	2.9	2.1	2.1	1.8	2.2	2.3	2.7	
Nitrate Bext	0.3	0.3	0.3	0.3	0.2	0.4	0.3	0.2	0.2	0.4	0.3	0.3	
POM Bext	1.1	0.1	0.8	1.7	1.9	0.3	0.5	0.1	0.1	0.2	0.1	0.1	
EC Bext	0.2	0.1	0.1	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	
Soil Bext	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	
Coarse Mass Bext	0.4	0.4	0.4	0.4	0.2	0.5	0.5	0.4	0.3	0.5	0.4	0.3	
Sea Salt Bext	0.9	1.6	0.7	0.7	0.7	1.0	1.3	1.4	1.4	1.7	1.8	1.4	




Craters of the Moon - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.0	13.6	7.5	14.3	14.9	14.0	12.8	12.8	12.8	16.2	14.5	13.6	11.0
Sulfate Bext	5.7	4.7	0.8	8.0	4.5	4.2	6.1	6.1	6.1	3.6	5.5	4.2	4.1
Nitrate Bext	11.3	8.6	1.0	16.0	13.8	6.9	8.6	8.6	10.3	7.6	9.2	7.3	8.6
POM Bext	9.1	11.9	4.0	4.3	10.7	14.8	6.5	6.5	5.4	22.9	13.9	13.0	4.3
EC Bext	1.9	2.1	0.4	1.4	2.1	2.5	1.7	1.7	1.8	3.3	2.7	1.9	1.0
Soil Bext	1.0	1.1	1.2	1.4	1.1	0.9	0.9	0.9	0.6	2.2	0.8	0.9	0.7
Coarse Mass Bext	2.9	3.4	4.0	2.8	2.7	3.5	2.9	2.9	2.8	7.1	2.3	3.0	1.8
Sea Salt Bext	0.0	0.1	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.1	0.1	0.0




Craters of the Moon - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.3	3.4	1.7	4.8	4.9	3.3	4.3	3.9	3.6	3.5	3.2	3.0	
Sulfate Bext	1.5	1.5	0.3	1.9	1.8	1.0	1.4	2.0	1.6	1.4	1.3	1.3	
Nitrate Bext	1.2	0.8	0.4	1.1	1.3	0.9	1.5	0.9	0.8	0.8	1.1	0.6	
POM Bext	1.3	0.8	0.5	1.5	1.4	0.9	1.4	0.9	0.7	0.8	0.7	0.7	
EC Bext	0.4	0.3	0.1	0.5	0.4	0.2	0.4	0.3	0.4	0.3	0.2	0.2	
Soil Bext	0.2	0.2	0.1	0.3	0.3	0.1	0.2	0.2	0.1	0.2	0.1	0.2	
Coarse Mass Bext	0.8	0.5	0.4	0.9	1.1	0.7	0.5	0.5	0.5	0.6	0.4	0.5	
Sea Salt Bext	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.1	0.1	0.0	0.0	




Hells Canyon - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.6	18.1	8.3	19.0	18.2			18.5	18.3	19.8	19.4	15.1	
Sulfate Bext	8.4	6.8	1.1	9.9	6.6			8.6	8.8	5.4	7.3	5.7	
Nitrate Bext	28.5	13.5	2.7	32.9	20.0			32.5	29.1	11.2	6.8	6.9	
POM Bext	15.6	31.4	3.7	11.4	20.7			14.7	13.1	43.9	51.0	17.8	
EC Bext	3.1	5.3	0.4	2.9	3.7			2.5	4.3	7.6	6.6	2.5	
Soil Bext	0.7	0.8	0.9	0.7	0.7			0.6	0.4	1.1	1.2	0.7	
Coarse Mass Bext	1.9	2.9	3.4	1.4	2.0			2.5	1.8	3.8	3.6	2.6	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0			0.1	0.2	0.2	0.1	0.0	




Hells Canyon - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.5	4.8	2.5	5.5	5.8			5.2	4.7	4.7	5.2	4.6	
Sulfate Bext	1.9	1.9	0.4	2.2	1.9			1.7	2.0	1.7	2.2	1.8	
Nitrate Bext	0.8	0.6	0.2	1.0	0.8			0.5	0.5	0.7	0.6	0.7	
POM Bext	2.0	1.5	0.5	1.6	2.3			2.1	1.4	1.5	1.7	1.4	
EC Bext	0.6	0.4	0.1	0.6	0.7			0.5	0.5	0.4	0.5	0.3	
Soil Bext	0.3	0.2	0.1	0.2	0.3			0.2	0.1	0.2	0.2	0.2	
Coarse Mass Bext	0.8	0.5	0.5	0.7	0.9			0.8	0.5	0.4	0.6	0.5	
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0			0.1	0.1	0.2	0.1	0.1	




Sawtooth - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.8	14.8	6.4	14.5	14.2	13.3	13.1	11.9	19.2	17.4			10.6
Sulfate Bext	3.1	3.5	0.8	3.5	2.1	2.6	4.0	3.8	3.8	4.1			2.5
Nitrate Bext	0.6	0.7	0.7	0.8	0.6	0.5	0.6	0.6	1.0	0.9			0.4
POM Bext	22.2	36.9	3.9	23.9	22.5	21.6	20.9	15.3	73.8	47.0			11.3
EC Bext	4.2	4.9	0.4	5.6	4.9	2.8	3.4	3.2	7.9	6.6			1.7
Soil Bext	0.8	1.0	0.9	1.1	0.8	0.5	0.8	0.7	1.1	1.4			0.8
Coarse Mass Bext	1.7	2.5	2.4	2.3	1.6	1.6	1.5	2.4	2.9	2.8			2.0
Sea Salt Bext	0.1	0.2	0.1	0.3	0.0	0.2	0.0	0.0	0.0	0.0			0.6




Sawtooth - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.0	3.8	1.5	4.9	3.8	3.2	4.1	3.5	4.0	3.8			3.8
Sulfate Bext	1.2	1.3	0.3	1.7	1.1	0.8	1.1	1.2	1.4	1.3			1.2
Nitrate Bext	0.3	0.2	0.3	0.4	0.3	0.2	0.3	0.2	0.3	0.2			0.2
POM Bext	2.4	2.1	0.6	2.8	2.1	1.9	2.6	1.9	2.1	2.3			2.2
EC Bext	0.6	0.5	0.1	0.9	0.6	0.5	0.6	0.5	0.7	0.5			0.4
Soil Bext	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.1	0.1	0.1			0.2
Coarse Mass Bext	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.4			0.3
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1			0.0




Sula Peak - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.4	17.0	7.4		12.6	12.1	16.9	12.0	14.6		24.9		11.6
Sulfate Bext	4.8	4.7	1.1	4.8	4.6	4.8	5.1	5.2			3.8		5.1
Nitrate Bext	1.5	1.7	1.1	1.3	1.6	1.6	1.6	1.4	1.4		2.7		1.0
POM Bext	20.0	59.5	4.8	11.5	12.6	43.3	12.7	30.6			136.5		11.3
EC Bext	2.5	8.8	0.4	2.0	2.1	4.3	1.7	4.9			19.8		1.7
Soil Bext	0.9	0.6	0.9	1.6	0.9	0.6	0.7	0.5			0.7		0.6
Coarse Mass Bext	2.5	3.6	2.7	3.7	2.4	1.7	2.1	2.8			5.2		2.8
Sea Salt Bext	0.3	0.0	0.3	1.0	0.0	0.0	0.0	0.0			0.1		0.0




Sula Peak - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.6	2.5	1.1	3.1	2.7	1.8	2.7	2.5	2.5	2.5	2.5	2.3	2.3
Sulfate Bext	1.1	1.3	0.3	1.3	1.1	0.7	1.2	1.4	1.3	1.3	1.3	1.2	1.2
Nitrate Bext	0.3	0.3	0.2	0.4	0.4	0.2	0.4	0.3	0.3	0.3	0.3	0.2	0.2
POM Bext	0.9	0.6	0.4	1.0	1.0	0.8	1.0	0.6	0.7	0.7	0.7	0.6	0.6
EC Bext	0.3	0.2	0.1	0.3	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.1	0.1
Soil Bext	0.1	0.1	0.1	0.2	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.3	0.2	0.3	0.2	0.1	0.2	0.3	0.4	0.4	0.4	0.4	0.4
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0




Yellowstone - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.8	11.5	6.4	12.9	12.6	11.4	11.2	10.8	9.6	11.8	12.6	14.0	9.6
Sulfate Bext	4.3	4.6	0.8	4.2	4.7	3.8	4.1	4.6	4.8	5.1	4.3	5.0	3.5
Nitrate Bext	1.8	1.6	0.6	2.0	2.4	1.6	1.4	1.5	1.6	0.9	2.4	1.7	1.6
POM Bext	13.5	15.5	4.6	18.4	16.1	11.2	12.8	8.9	6.7	15.1	14.7	25.6	15.2
EC Bext	2.5	2.2	0.4	3.7	2.6	1.9	2.3	1.9	1.8	2.3	2.4	2.6	1.7
Soil Bext	1.0	0.8	1.0	0.8	1.4	1.2	0.4	0.9	0.6	1.1	0.8	1.0	0.6
Coarse Mass Bext	2.6	2.4	3.0	2.8	2.7	3.1	1.3	2.9	1.8	2.6	2.6	2.6	2.3
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.1	0.1




Yellowstone - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.6	2.0	0.4	2.8	3.0	2.6	1.9	2.6	2.1	2.3	2.2	1.8	1.9
Sulfate Bext	1.5	1.4	0.3	1.7	1.9	1.3	1.1	1.3	1.5	1.6	1.2	1.1	1.4
Nitrate Bext	0.7	0.5	0.3	0.9	0.8	0.8	0.5	0.6	0.3	0.4	0.7	0.6	0.6
POM Bext	1.1	0.8	0.5	1.1	1.2	1.1	0.9	1.3	1.1	0.8	1.0	0.8	0.5
EC Bext	0.3	0.3	0.1	0.2	0.4	0.3	0.2	0.5	0.3	0.4	0.3	0.1	0.2
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.2	0.2	0.3	0.2	0.3	0.2	0.3	0.1	0.2	0.2	0.3	0.3
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Mammoth Cave - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	31.4	30.7	11.1	32.1	31.5	31.0	31.8	30.4	33.7	31.9	32.2	28.7	26.9
Sulfate Bext	188.9	181.9	4.5	3	3	4	6	0	7	7	1	0	9
Nitrate Bext	9.7	9.2	0.8	7.8	11.6	10.1	9.7	9.5	8.5	12.5	7.2	6.2	11.8
POM Bext	22.3	17.7	10.7	30.3	20.1	21.6	20.1	19.2	17.5	19.1	25.9	14.4	11.7
EC Bext	6.2	5.7	0.3	8.7	5.5	5.6	6.1	5.2	7.6	6.5	6.0	4.4	4.0
Soil Bext	0.9	0.8	1.0	0.6	0.9	1.3	0.8	1.1	0.6	0.8	0.9	1.0	0.7
Coarse Mass Bext	2.2	3.0	2.3	3.6	2.3	1.1	2.1	1.8	2.6	3.1	3.9	3.0	2.4
Sea Salt Bext	0.1	0.2	0.2	0.2	0.0	0.0	0.0	0.3	0.2	0.3	0.2	0.0	0.1




Mammoth Cave - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	16.5	15.8	5.0	17.1	15.2	17.4	16.7	16.2	17.3	16.6	15.7	15.1	14.1
Sulfate Bext	22.1	20.9	0.9	25.4	19.2	25.6	20.1	20.3	25.3	23.2	19.8	19.3	17.0
Nitrate Bext	8.8	7.4	0.7	7.1	8.6	9.5	10.1	8.6	9.1	7.3	8.9	6.7	5.2
POM Bext	6.3	5.2	2.5	6.9	4.5	6.8	7.1	6.2	5.8	6.3	4.5	5.1	4.3
EC Bext	3.0	2.7	0.1	3.7	2.4	3.0	3.1	2.9	3.5	3.3	2.3	2.3	1.9
Soil Bext	0.3	0.3	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Coarse Mass Bext	1.4	1.7	0.9	1.5	0.9	1.1	1.8	1.5	1.9	1.9	2.1	1.4	1.5
Sea Salt Bext	0.1	0.3	0.1	0.0	0.0	0.0	0.0	0.4	0.2	0.4	0.4	0.3	0.1




Acadia - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	22.9	21.5	12.4	21.6	23.3	23.9	23.6	22.0	23.0	23.4	21.7	20.2	18.9
Sulfate Bext	69.2	57.2	4.8	48.4	75.5	74.8	84.9	62.5	70.1	69.6	56.4	46.7	43.5
Nitrate Bext	8.0	6.4	2.0	10.6	7.5	8.2	5.6	8.3	5.7	9.2	7.0	5.5	4.8
POM Bext	11.2	9.6	11.8	8.2	11.9	14.6	12.8	8.3	13.0	9.5	11.1	8.2	6.1
EC Bext	4.3	3.5	0.4	4.6	4.5	4.0	4.9	3.6	4.6	4.1	3.9	2.8	2.4
Soil Bext	0.5	0.3	0.5	0.3	0.6	0.8	0.5	0.4	0.3	0.4	0.4	0.3	0.4
Coarse Mass Bext	1.9	2.3	2.2	2.5	1.6	1.9	1.7	1.8	2.0	2.8	2.7	2.1	2.0
Sea Salt Bext	1.3	1.6	2.1	3.1	0.0	0.8	0.6	1.9	1.3	1.3	1.9	1.3	2.0




Acadia - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	8.8	7.7	4.7	8.9	8.9	8.8	8.8	8.6	7.6	8.2	8.2	7.8	6.9
Sulfate Bext	6.8	5.0	0.8	7.2	7.6	6.3	6.8	5.9	4.7	5.7	5.9	4.9	3.9
Nitrate Bext	1.1	0.7	0.3	1.0	1.4	1.0	1.1	0.9	0.9	0.9	0.7	0.7	0.5
POM Bext	2.2	1.9	2.0	2.2	1.8	2.5	2.1	2.6	1.8	1.9	2.3	2.0	1.5
EC Bext	0.9	0.7	0.1	1.1	0.9	0.9	0.8	0.8	0.8	0.9	0.7	0.6	0.5
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.7	0.8	0.6	0.7	0.6	0.8	0.6	0.9	0.7	0.8	0.7	0.9	1.0
Sea Salt Bext	0.4	0.6	0.2	0.3	0.1	0.6	0.7	0.5	0.8	0.6	0.4	0.8	0.6




Moosehorn - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	21.7	19.9	12.0	20.6	22.1	23.1	22.5	20.3	22.4	21.6	19.2	18.7	17.7
Sulfate Bext	58.5	45.7	4.0	43.5	66.1	71.5	64.7	46.5	61.2	55.8	37.9	37.8	35.8
Nitrate Bext	6.4	4.6	1.7	8.5	5.9	6.4	5.5	5.7	4.3	6.1	5.2	3.9	3.5
POM Bext	11.9	9.7	11.6	8.8	12.8	16.0	13.5	8.7	14.5	9.6	9.3	8.7	6.3
EC Bext	4.4	3.1	0.4	3.6	4.0	4.0	7.3	3.2	3.9	3.3	3.1	2.5	2.9
Soil Bext	0.4	0.3	0.4	0.3	0.5	0.4	0.4	0.4	0.3	0.3	0.3	0.4	0.4
Coarse Mass Bext	2.1	1.7	2.3	2.7	2.1	1.7	2.3	1.8	1.6	1.6	1.9	1.9	1.5
Sea Salt Bext	0.9	1.8	1.5	2.1	0.0	0.5	0.4	1.8	2.6	1.4	1.4	1.2	2.3




Moosehorn - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	9.2	7.8	5.0	8.9	9.3	9.1	9.5	8.9	8.0	8.6	7.8	7.8	6.8
Sulfate Bext	6.7	5.0	0.8	6.6	7.0	6.6	7.4	5.8	5.1	5.5	5.5	4.8	4.0
Nitrate Bext	1.1	0.7	0.3	1.2	1.3	1.0	0.9	0.9	0.7	0.9	0.6	0.6	0.5
POM Bext	3.1	2.1	2.2	3.0	3.0	3.1	3.1	3.1	2.2	2.9	1.9	2.0	1.6
EC Bext	1.0	0.7	0.1	1.2	1.0	1.0	0.9	0.9	0.9	1.1	0.6	0.6	0.4
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	1.1	0.8	0.8	1.0	0.9	1.1	1.2	1.1	0.7	0.7	0.8	1.0	0.6
Sea Salt Bext	0.3	0.6	0.2	0.0	0.1	0.3	0.3	0.7	0.8	0.7	0.3	0.7	0.6




Isle Royale - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	20.7	21.8	12.4	20.2	22.5	21.5	19.9	19.6	23.5	21.8	21.7	21.1	20.6
Sulfate Bext	33.6	43.8	3.6	31.5	40.8	32.5	29.5	33.4	57.6	45.9	46.7	31.7	37.1
Nitrate Bext	19.8	20.9	2.2	16.9	32.6	17.4	18.3	13.6	19.3	17.6	15.9	31.8	19.7
POM Bext	11.9	10.8	13.3	8.4	8.5	20.0	11.4	11.0	12.9	12.0	11.2	10.6	7.3
EC Bext	3.6	3.5	0.4	3.9	3.9	4.2	3.4	2.9	4.7	3.8	3.4	3.0	2.7
Soil Bext	0.4	0.5	0.4	0.4	0.4	0.5	0.4	0.5	0.5	0.5	0.6	0.4	0.5
Coarse Mass Bext	3.4	3.5	3.5	4.9	2.1	3.9	2.7	3.5	3.6	4.1	3.9	2.8	3.4
Sea Salt Bext	0.2	0.2	0.4	0.0	0.5	0.0	0.0	0.3	0.3	0.3	0.2	0.2	0.1




Isle Royale - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.8	6.1	3.7	6.5	7.2	7.1	7.0	6.1	7.1	6.4	6.4	5.6	5.1
Sulfate Bext	4.6	3.8	0.7	4.4	5.4	5.4	4.2	3.7	5.2	4.3	4.0	3.2	2.5
Nitrate Bext	0.7	0.4	0.1	0.6	0.9	0.5	0.9	0.5	0.5	0.6	0.5	0.3	0.2
POM Bext	1.2	1.0	1.0	1.0	1.2	1.1	1.6	1.1	1.2	0.8	1.0	1.1	1.0
EC Bext	0.4	0.3	0.1	0.4	0.3	0.4	0.6	0.3	0.4	0.3	0.4	0.2	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.7	0.7	0.5	0.6	0.6	0.8	0.8	0.7	0.8	0.7	0.8	0.6	0.7
Sea Salt Bext	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.2	0.3	0.2	0.1	0.1




Seney - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	24.2	24.2	12.7	23.0	25.6	24.6	24.5	23.2	26.2	24.8	26.6	22.2	21.3
Sulfate Bext	60.7	66.5	4.7	49.6	66.8	63.6	62.9	60.6	90.6	67.5	91.3	43.8	39.2
Nitrate Bext	26.9	24.3	2.3	22.0	40.0	27.0	28.1	17.5	19.1	29.5	21.7	27.1	24.2
POM Bext	13.3	13.7	14.1	11.8	12.5	12.5	16.0	13.4	18.2	12.0	21.0	9.2	8.1
EC Bext	4.3	4.4	0.5	4.8	4.6	3.9	4.7	3.5	5.9	5.1	4.7	3.3	2.8
Soil Bext	0.5	0.5	0.5	0.4	0.5	0.6	0.4	0.6	0.5	0.6	0.7	0.5	0.4
Coarse Mass Bext	1.9	2.1	2.2	1.8	1.5	2.6	2.1	1.5	2.4	2.5	2.3	1.7	1.4
Sea Salt Bext	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.2	0.1	0.1




Seney - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.1	6.5	3.7	6.5	6.8	7.8	8.0	6.6	7.5	6.7	6.8	5.9	5.3
Sulfate Bext	4.8	4.3	0.6	3.4	4.5	6.3	5.8	4.0	5.7	4.6	4.7	3.8	2.7
Nitrate Bext	0.8	0.5	0.1	0.6	0.9	0.8	1.1	0.6	0.6	0.8	0.6	0.4	0.3
POM Bext	1.6	1.2	1.1	1.6	1.2	1.6	2.1	1.5	1.7	1.0	1.3	1.1	0.9
EC Bext	0.5	0.4	0.1	0.5	0.4	0.5	0.6	0.4	0.6	0.4	0.4	0.2	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.7	0.6	0.5	1.1	0.5	0.6	0.5	0.6	0.6	0.5	0.6	0.5	0.7
Sea Salt Bext	0.1	0.2	0.1	0.0	0.1	0.1	0.2	0.2	0.2	0.3	0.3	0.2	0.2




Boundary Waters - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	19.6	20.1	11.6	19.9	20.0	20.1	20.1	17.8	21.3	19.6	19.8	19.7	20.0
Sulfate Bext	27.4	33.3	3.1	27.0	27.3	31.7	27.4	23.8	50.4	30.0	33.4	24.1	28.7
Nitrate Bext	17.4	18.6	1.8	16.4	25.0	12.1	20.3	13.3	14.3	17.8	14.8	20.6	25.5
POM Bext	12.4	10.4	13.4	11.5	8.3	17.6	14.6	10.1	8.9	11.9	10.4	13.4	7.4
EC Bext	3.1	2.9	0.4	4.4	3.0	3.3	2.6	2.2	3.5	3.2	3.0	2.7	2.3
Soil Bext	0.5	0.5	0.6	0.8	0.4	0.6	0.4	0.4	0.5	0.4	0.6	0.5	0.5
Coarse Mass Bext	2.6	2.1	3.1	3.7	2.1	2.5	2.2	2.2	2.5	1.9	1.5	1.7	2.8
Sea Salt Bext	0.1	0.2	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.2	0.2	0.6	0.1




Boundary Waters - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.4	5.6	3.4	6.0	6.9	7.0	6.5	5.8	6.3	5.7	5.6	5.5	4.6
Sulfate Bext	4.1	3.8	0.7	3.8	4.8	4.9	3.6	3.4	4.6	4.1	3.4	3.9	2.9
Nitrate Bext	0.7	0.4	0.2	0.7	0.8	0.7	0.8	0.5	0.6	0.4	0.4	0.3	0.3
POM Bext	1.9	1.3	1.5	1.6	1.8	2.1	2.5	1.6	1.6	1.3	1.6	1.2	0.9
EC Bext	0.6	0.4	0.1	0.6	0.4	0.7	0.7	0.4	0.4	0.5	0.4	0.3	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.7	0.5	0.5	0.5	0.7	0.6	0.8	0.7	0.5	0.5	0.5	0.6	0.5
Sea Salt Bext	0.2	0.1	0.1	0.0	0.6	0.0	0.0	0.1	0.1	0.1	0.3	0.1	0.1




Voyageurs - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	19.3	19.8	12.1	19.1	18.6	20.1	20.2	18.4	19.9	20.5	19.2	19.2	20.2
Sulfate Bext	21.3	28.9	2.7	22.5	18.2	23.6	24.4	18.0	41.6	29.8	27.4	18.6	27.1
Nitrate Bext	18.2	18.4	1.5	14.6	22.5	15.7	21.5	16.5	11.8	15.4	18.2	21.4	25.1
POM Bext	13.6	11.7	14.1	10.0	7.7	22.1	14.3	13.8	8.9	18.9	9.8	12.7	8.3
EC Bext	3.2	3.2	0.4	3.7	2.8	3.7	3.3	2.6	3.5	3.9	2.9	2.9	2.7
Soil Bext	0.4	0.5	0.5	0.5	0.3	0.5	0.5	0.4	0.4	0.6	0.5	0.4	0.5
Coarse Mass Bext	2.8	2.5	3.5	3.8	2.1	2.9	2.6	2.6	2.5	3.5	1.7	2.1	2.7
Sea Salt Bext	0.5	0.3	0.6	1.6	0.3	0.2	0.1	0.3	0.2	0.3	0.3	0.7	0.2




Voyageurs - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.1	6.4	4.3	7.0	7.1	7.5	7.7	6.4	6.8	6.5	6.6	6.5	5.5
Sulfate Bext	4.2	3.7	0.8	4.4	4.5	4.6	4.4	3.1	4.1	3.9	3.7	3.8	3.0
Nitrate Bext	0.8	0.5	0.1	1.0	0.8	0.8	0.8	0.5	0.5	0.6	0.5	0.4	0.3
POM Bext	2.1	1.6	1.6	1.6	2.0	2.3	2.8	1.9	1.8	1.5	2.2	1.7	1.1
EC Bext	0.6	0.4	0.1	0.7	0.4	0.6	0.7	0.5	0.5	0.6	0.4	0.3	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.7	0.6	0.5	0.4	0.7	0.7	0.9	0.8	0.8	0.5	0.5	0.8	0.5
Sea Salt Bext	0.1	0.1	0.1	0.1	0.1	0.2	0.0	0.2	0.2	0.1	0.3	0.1	0.1




Hercules-Glade - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	26.7	26.0	11.3			27.2	26.6	26.5	29.5	25.9	26.2	25.0	23.6
Sulfate Bext	87.9	86.7	2.9			104.2	87.0	72.6	139.2	76.8	92.6	73.0	51.9
Nitrate Bext	17.9	17.4	1.0			14.6	15.0	24.1	15.3	20.3	10.0	19.4	22.0
POM Bext	25.3	18.5	13.1			18.2	27.4	30.3	22.8	19.5	21.9	13.8	14.3
EC Bext	5.2	5.1	0.3			5.0	5.6	5.1	6.7	5.6	5.6	3.6	4.0
Soil Bext	0.9	0.9	1.0			1.0	0.8	0.9	0.7	1.5	0.8	0.8	0.7
Coarse Mass Bext	2.8	3.5	2.7			2.9	2.5	2.9	4.1	3.3	3.9	3.1	3.2
Sea Salt Bext	0.2	0.5	0.2			0.0	0.0	0.6	0.8	0.6	0.4	0.2	0.3




Hercules-Glade - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.8	12.5	4.7			13.6	12.2	12.7	13.6	13.3	13.0	11.7	11.0
Sulfate Bext	10.8	11.8	0.7			11.7	9.3	11.6	14.8	13.7	10.2	10.5	9.5
Nitrate Bext	4.9	3.9	0.5			6.1	4.3	4.3	4.2	3.9	6.0	2.8	2.4
POM Bext	5.1	4.5	2.3			5.5	5.3	4.6	4.8	4.9	5.1	4.1	3.7
EC Bext	2.0	1.9	0.1			2.4	1.8	1.8	2.2	2.5	1.9	1.4	1.4
Soil Bext	0.3	0.3	0.2			0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.3
Coarse Mass Bext	2.3	2.3	1.1			2.4	2.3	2.1	2.8	1.8	2.5	2.5	2.0
Sea Salt Bext	0.2	0.1	0.1			0.0	0.2	0.3	0.1	0.2	0.2	0.1	0.2




Mingo - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	28.4	27.1	11.6		29.5	29.6	28.9	25.7		27.3	29.1	26.8	25.1
Sulfate Bext	104.6	93.7	2.4		91.6	141.3	113.3	72.1		101.8	120.9	87.9	64.0
Nitrate Bext	27.2	12.5	1.6		39.3	18.4	29.7	21.5		9.4	10.7	14.4	15.3
POM Bext	20.5	22.6	11.5		38.5	13.0	17.5	13.2		17.0	32.3	20.1	20.7
EC Bext	5.5	6.4	0.4		9.0	3.8	5.2	4.1		6.2	8.1	5.6	5.8
Soil Bext	1.4	2.0	0.9		1.6	1.7	1.1	1.2		2.7	2.1	1.7	1.4
Coarse Mass Bext	6.5	7.4	3.2		8.4	5.1	4.5	7.8		9.2	8.4	5.2	6.7
Sea Salt Bext	0.2	0.3	0.5		0.0	0.1	0.0	0.7		0.4	0.4	0.3	0.3




Mingo - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.3	13.9	5.4	13.7	15.2	14.7	13.7		14.2	15.1	13.9	12.3	
Sulfate Bext	13.8	13.6	0.9	13.2	15.8	14.0	12.5		14.1	14.5	14.0	11.8	
Nitrate Bext	4.4	4.2	0.4	5.1	4.3	4.0	4.4		4.1	6.1	4.5	2.1	
POM Bext	6.3	5.0	2.6	5.0	7.5	7.8	4.8		5.0	5.8	5.1	4.2	
EC Bext	2.3	2.1	0.1	2.1	2.6	2.7	1.7		2.4	2.6	1.8	1.6	
Soil Bext	0.4	0.5	0.2	0.4	0.6	0.5	0.3		0.6	0.7	0.5	0.3	
Coarse Mass Bext	3.2	3.2	0.9	2.1	3.2	3.2	4.1		3.5	4.1	3.0	2.4	
Sea Salt Bext	0.1	0.2	0.1	0.0	0.0	0.0	0.3		0.4	0.3	0.2	0.1	




Cabinet Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.1	13.5	7.5	13.7	13.9	14.6	14.2	12.6	15.4	14.8	12.5	12.0	
Sulfate Bext	6.5	7.1	1.1	7.0	5.9	6.3	6.7	8.4	7.0	6.5	6.9	6.8	
Nitrate Bext	2.0	1.6	1.1	1.4	2.8	2.5	1.3	1.5	2.0	1.4	1.4	1.6	
POM Bext	16.9	15.9	4.3	14.8	16.0	18.9	18.1	9.6	25.5	24.2	10.6	9.8	
EC Bext	2.8	2.8	0.4	2.5	3.0	3.0	2.7	2.7	4.1	3.7	2.0	1.7	
Soil Bext	1.0	1.0	1.0	1.7	1.1	0.6	0.7	0.8	1.0	0.9	1.3	1.0	
Coarse Mass Bext	2.8	2.9	3.4	4.3	2.5	2.2	2.3	2.5	2.5	3.0	3.5	2.8	
Sea Salt Bext	0.1	0.1	0.2	0.3	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	




Cabinet Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.6	3.2	1.5	4.3	3.7	3.3	3.2	4.0	2.4	3.6	3.2	2.7	
Sulfate Bext	1.8	1.7	0.4	2.2	2.0	1.5	1.5	2.5	1.2	1.7	1.5	1.6	
Nitrate Bext	0.8	0.6	0.4	0.8	0.7	0.8	0.7	0.6	0.4	0.6	0.7	0.5	
POM Bext	1.0	0.8	0.4	1.3	1.0	0.9	0.9	0.9	0.5	1.2	0.9	0.5	
EC Bext	0.3	0.3	0.1	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.2
Soil Bext	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.3	0.3	0.2	0.4	0.4	0.2	0.3	0.4	0.2	0.3	0.2	0.2	
Sea Salt Bext	0.1	0.1	0.0	0.1	0.2	0.2	0.1	0.3	0.1	0.1	0.1	0.1	




Gates of the Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.3	11.2	6.4	10.8	10.6	12.8	10.9	11.3	12.7		11.0	9.9	
Sulfate Bext	5.4	6.0	1.1	6.0	5.1	5.1	5.5	6.6	6.6		5.4	5.4	
Nitrate Bext	1.9	1.6	1.0	1.9	1.8	2.3	1.5	2.0	2.7		0.8	1.0	
POM Bext	11.3	11.8	5.0	8.7	9.8	16.2	10.3	10.7	17.2		11.7	7.7	
EC Bext	1.8	1.9	0.5	1.6	1.6	2.5	1.6	2.1	2.7		1.6	1.2	
Soil Bext	0.7	0.8	0.8	1.1	0.6	0.5	0.8	0.6	1.0		0.8	0.7	
Coarse Mass Bext	1.7	1.8	1.8	1.9	1.4	1.6	1.8	1.4	1.8		1.9	2.0	
Sea Salt Bext	0.1	0.0	0.1	0.0	0.0	0.0	0.2	0.0	0.0		0.0	0.0	




Gates of the Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.7	0.9	0.3		1.8	2.2	0.9	1.9	1.1	0.9		0.8	0.8
Sulfate Bext	1.0	0.9	0.2		1.2	1.2	0.6	0.9	1.0	0.8		0.8	1.0
Nitrate Bext	0.5	0.3	0.3		0.5	0.6	0.3	0.5	0.3	0.3		0.3	0.2
POM Bext	0.7	0.4	0.4		0.6	0.8	0.5	0.9	0.5	0.3		0.5	0.3
EC Bext	0.2	0.1	0.1		0.2	0.2	0.1	0.2	0.2	0.2		0.1	0.1
Soil Bext	0.1	0.1	0.1		0.1	0.1	0.1	0.1	0.1	0.1		0.1	0.1
Coarse Mass Bext	0.4	0.2	0.3		0.4	0.5	0.3	0.4	0.2	0.2		0.2	0.2
Sea Salt Bext	0.0	0.0	0.0		0.0	0.1	0.0	0.0	0.0	0.1		0.0	0.0




Glacier - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	22.3	18.7	9.2	21.3	20.4	19.7	30.2	19.7	19.1	18.6	21.2	16.0	
Sulfate Bext	11.4	12.0	1.3	12.7	9.9	13.4	9.7	11.2	16.3	10.3	10.2	11.2	
Nitrate Bext	9.4	7.3	2.1	9.5	9.1	8.9	7.3	12.1	11.9	6.0	6.2	5.2	
POM Bext	87.7	28.6	5.0	38.6	46.9	32.0	292.8	28.1	22.2	25.8	52.5	14.1	
EC Bext	11.2	5.7	0.4	7.5	7.4	6.6	29.2	5.3	6.5	6.1	6.7	3.7	
Soil Bext	1.4	1.1	1.3	1.9	1.3	1.2	1.4	1.0	0.6	1.3	1.5	1.0	
Coarse Mass Bext	5.2	4.6	4.6	7.4	5.2	2.5	6.5	4.6	2.2	5.3	6.9	3.9	
Sea Salt Bext	0.3	0.2	0.3	0.2	0.8	0.0	0.1	0.2	0.1	0.2	0.2	0.2	




Glacier - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.2	7.0	2.4	7.5	7.0	6.9	7.6	7.1	8.3	6.1	6.9	6.8	
Sulfate Bext	3.0	3.1	0.4	3.4	3.1	3.0	2.9	2.8	4.2	2.2	2.8	3.3	
Nitrate Bext	0.9	0.8	0.3	0.9	0.9	0.8	1.0	0.7	0.8	0.7	0.9	0.9	
POM Bext	3.3	2.9	0.5	3.5	3.1	2.7	3.8	3.4	3.7	2.5	2.9	2.6	
EC Bext	1.4	1.5	0.1	1.4	1.2	1.3	1.9	1.3	1.9	1.3	1.5	1.1	
Soil Bext	0.2	0.2	0.1	0.1	0.2	0.3	0.2	0.2	0.2	0.1	0.2	0.2	
Coarse Mass Bext	0.8	0.8	0.3	0.8	0.7	0.9	0.6	1.2	1.2	0.5	0.7	0.6	
Sea Salt Bext	0.1	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.2	0.1	0.1	




Medicine Lake - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.7	18.0	7.9	17.8	18.3	16.6	16.9	19.0	17.7	18.8	18.7	18.2	16.7
Sulfate Bext	17.0	19.1	1.1	14.9	18.1	15.9	14.1	21.8	19.4	19.9	22.2	17.4	16.7
Nitrate Bext	16.3	15.8	1.0	13.4	19.1	13.0	13.3	22.5	15.2	18.9	12.7	16.1	15.9
POM Bext	9.5	9.6	4.0	12.3	8.0	6.9	9.4	10.9	7.0	12.5	12.1	11.7	4.8
EC Bext	2.3	2.2	0.4	3.4	2.2	2.0	2.2	1.9	2.5	2.5	2.4	1.9	1.5
Soil Bext	0.8	0.7	1.0	1.1	1.0	0.7	0.5	0.5	0.6	0.7	0.9	0.8	0.7
Coarse Mass Bext	4.5	4.4	3.9	6.2	5.5	3.8	4.2	2.5	4.3	3.7	7.1	4.0	3.0
Sea Salt Bext	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.2	0.1	0.1	0.1	1.9	0.8




Medicine Lake - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.3	6.5	3.0	8.0	7.6	6.8	6.9	7.0	7.0	5.8	6.5	6.9	6.4
Sulfate Bext	3.5	3.5	0.4	4.3	3.4	3.1	3.3	3.4	4.2	3.0	3.2	3.6	3.7
Nitrate Bext	1.2	1.1	0.4	1.8	1.1	0.8	1.0	1.2	1.2	1.0	1.0	1.2	0.9
POM Bext	1.9	1.3	0.7	2.3	2.0	1.7	1.8	2.0	1.5	1.0	1.3	1.5	1.1
EC Bext	0.6	0.5	0.1	0.8	0.7	0.6	0.5	0.5	0.7	0.5	0.6	0.5	0.4
Soil Bext	0.4	0.3	0.2	0.3	0.5	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.3
Coarse Mass Bext	2.2	1.6	0.7	1.8	2.9	2.2	2.1	1.8	1.4	1.3	1.8	1.8	1.8
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0




Monture - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.5	15.3	7.7	15.4	13.9	15.2	13.4	15.6	14.6	21.9	12.5	12.0	
Sulfate Bext	5.1	6.2	1.0	5.3	5.1	4.8	5.3	7.3	6.2	5.8	5.7	6.1	
Nitrate Bext	1.4	1.3	1.2	1.4	1.6	1.6	1.1	2.0	1.0	1.3	1.5	0.7	
POM Bext	22.3	30.9	4.5	21.7	19.6	30.5	17.4	26.5	22.4	81.6	13.1	10.9	
EC Bext	2.8	4.0	0.4	3.0	2.6	3.3	2.3	5.0	3.6	8.2	1.7	1.6	
Soil Bext	1.3	1.5	1.2	2.0	1.5	0.8	0.8	1.1	1.4	2.4	1.1	1.4	
Coarse Mass Bext	3.6	2.8	3.9	5.6	4.0	2.4	2.4	2.6	2.7	3.8	1.7	3.1	
Sea Salt Bext	0.0	0.2	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.6	0.0	




Monture - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.9	3.1	1.5	4.9	3.5	3.1	3.9	3.8	3.1	2.8	2.8	3.1	
Sulfate Bext	1.5	1.5	0.3	2.1	1.5	1.0	1.4	2.0	1.1	1.5	1.2	1.7	
Nitrate Bext	0.5	0.3	0.3	0.8	0.5	0.4	0.3	0.3	0.3	0.3	0.3	0.2	0.3
POM Bext	1.8	1.2	0.5	2.3	1.4	1.5	2.1	1.4	1.3	0.8	1.1	1.1	
EC Bext	0.5	0.4	0.1	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.3	0.3
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.3	0.2	0.2	0.5	0.3	0.1	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Sea Salt Bext	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.2	0.1	0.1	0.0	0.0




Sula Peak - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.4	17.0	7.4		12.6	12.1	16.9	12.0	14.6		24.9		11.6
Sulfate Bext	4.8	4.7	1.1	4.8	4.6	4.8	5.1	5.2			3.8		5.1
Nitrate Bext	1.5	1.7	1.1	1.3	1.6	1.6	1.6	1.4	1.4		2.7		1.0
POM Bext	20.0	59.5	4.8	11.5	12.6	43.3	12.7	30.6			136.5		11.3
EC Bext	2.5	8.8	0.4	2.0	2.1	4.3	1.7	4.9			19.8		1.7
Soil Bext	0.9	0.6	0.9	1.6	0.9	0.6	0.7	0.5			0.7		0.6
Coarse Mass Bext	2.5	3.6	2.7	3.7	2.4	1.7	2.1	2.8			5.2		2.8
Sea Salt Bext	0.3	0.0	0.3	1.0	0.0	0.0	0.0	0.0			0.1		0.0




Sula Peak - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.6	2.5	1.1	3.1	2.7	1.8	2.7	2.5		2.5			2.3
Sulfate Bext	1.1	1.3	0.3	1.3	1.1	0.7	1.2	1.4		1.3			1.2
Nitrate Bext	0.3	0.3	0.2	0.4	0.4	0.2	0.4	0.3		0.3			0.2
POM Bext	0.9	0.6	0.4	1.0	1.0	0.8	1.0	0.6		0.7			0.6
EC Bext	0.3	0.2	0.1	0.3	0.3	0.2	0.3	0.2		0.2			0.1
Soil Bext	0.1	0.1	0.1	0.2	0.1	0.0	0.1	0.1		0.1			0.1
Coarse Mass Bext	0.2	0.3	0.2	0.3	0.2	0.1	0.2	0.3		0.4			0.4
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0		0.1			0.0




UL Bend - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.1	14.9	8.2	15.6	13.6	17.1	14.2	13.9	15.6	16.6	14.4	13.8	
Sulfate Bext	9.8	10.3	1.0	9.6	11.2	8.5	9.8	10.9	11.2	9.8	8.8	10.8	
Nitrate Bext	8.0	5.7	1.5	13.3	6.1	8.3	4.3	4.7	6.4	5.3	5.8	6.3	
POM Bext	12.8	13.3	4.5	11.6	6.2	19.7	13.6	8.6	13.9	26.2	10.7	7.2	
EC Bext	2.1	2.3	0.4	2.0	1.5	3.1	1.7	2.7	2.4	3.7	1.5	1.2	
Soil Bext	0.8	0.9	1.0	0.8	0.5	1.1	0.7	0.6	1.1	0.9	0.9	0.8	
Coarse Mass Bext	4.0	4.0	3.9	3.7	2.9	5.9	3.4	3.7	4.7	4.5	3.7	3.6	
Sea Salt Bext	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.2	0.1	




UL Bend - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.7	4.3	2.5	5.2	4.6	4.6	4.6	4.4	4.0	4.4	4.3	4.6	
Sulfate Bext	1.8	1.8	0.3	2.1	1.6	1.6	1.8	1.8	1.4	1.8	1.9	2.0	
Nitrate Bext	0.5	0.3	0.2	0.6	0.5	0.4	0.5	0.3	0.2	0.4	0.5	0.3	
POM Bext	1.2	1.0	0.5	1.3	1.1	1.3	1.2	1.2	1.0	0.8	0.9	0.9	
EC Bext	0.5	0.3	0.1	0.6	0.5	0.5	0.4	0.4	0.5	0.3	0.3	0.3	
Soil Bext	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.1	0.2	0.2	
Coarse Mass Bext	0.9	0.9	0.6	1.1	1.0	0.8	0.8	0.8	0.7	1.1	0.7	1.0	
Sea Salt Bext	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	




Yellowstone - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.8	11.5	6.4	12.9	12.6	11.4	11.2	10.8	9.6	11.8	12.6	14.0	9.6
Sulfate Bext	4.3	4.6	0.8	4.2	4.7	3.8	4.1	4.6	4.8	5.1	4.3	5.0	3.5
Nitrate Bext	1.8	1.6	0.6	2.0	2.4	1.6	1.4	1.5	1.6	0.9	2.4	1.7	1.6
POM Bext	13.5	15.5	4.6	18.4	16.1	11.2	12.8	8.9	6.7	15.1	14.7	25.6	15.2
EC Bext	2.5	2.2	0.4	3.7	2.6	1.9	2.3	1.9	1.8	2.3	2.4	2.6	1.7
Soil Bext	1.0	0.8	1.0	0.8	1.4	1.2	0.4	0.9	0.6	1.1	0.8	1.0	0.6
Coarse Mass Bext	2.6	2.4	3.0	2.8	2.7	3.1	1.3	2.9	1.8	2.6	2.6	2.6	2.3
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.1	0.1




Yellowstone - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.6	2.0	0.4	2.8	3.0	2.6	1.9	2.6	2.1	2.3	2.2	1.8	1.9
Sulfate Bext	1.5	1.4	0.3	1.7	1.9	1.3	1.1	1.3	1.5	1.6	1.2	1.1	1.4
Nitrate Bext	0.7	0.5	0.3	0.9	0.8	0.8	0.5	0.6	0.3	0.4	0.7	0.6	0.6
POM Bext	1.1	0.8	0.5	1.1	1.2	1.1	0.9	1.3	1.1	0.8	1.0	0.8	0.5
EC Bext	0.3	0.3	0.1	0.2	0.4	0.3	0.2	0.5	0.3	0.4	0.3	0.1	0.2
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.2	0.2	0.3	0.2	0.3	0.2	0.3	0.1	0.2	0.2	0.3	0.3
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.1	12.4	7.9	12.7	12.8	12.1	10.7	11.3	13.8	13.4	12.5	11.1	
Sulfate Bext	4.0	4.4	0.8	4.0	3.2	4.1	4.7	5.4	4.7	4.0	3.7	4.4	
Nitrate Bext	1.1	1.4	0.9	1.0	1.5	0.7	1.1	2.7	1.0	1.7	1.1	0.5	
POM Bext	10.0	10.0	4.2	10.6	13.4	9.7	6.5	6.3	13.9	10.4	12.1	7.3	
EC Bext	1.6	1.7	0.4	2.3	1.8	1.5	1.0	1.2	3.2	1.6	1.5	1.1	
Soil Bext	2.4	2.6	1.3	2.8	3.0	1.8	1.9	1.6	2.5	3.7	2.6	2.3	
Coarse Mass Bext	5.5	5.9	4.3	6.6	4.7	5.9	4.6	4.6	6.5	7.8	4.8	5.7	
Sea Salt Bext	0.1	0.2	0.4	0.1	0.0	0.0	0.1	0.3	0.0	0.1	0.0	0.3	




Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.6	2.2	1.1	2.7	2.4	2.5	2.5	2.5	2.5	2.4	2.2	1.9	1.8
Sulfate Bext	1.2	1.1	0.3	1.5	1.1	1.0	1.3	1.3	1.3	1.2	1.1	1.1	1.0
Nitrate Bext	0.3	0.2	0.2	0.4	0.2	0.3	0.3	0.3	0.2	0.1	0.3	0.2	0.2
POM Bext	0.8	0.5	0.4	0.7	0.7	0.9	0.7	0.7	0.8	0.6	0.5	0.2	0.3
EC Bext	0.3	0.2	0.1	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.1	0.1
Soil Bext	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.3	0.3	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.4	0.3	0.3	0.3
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.0




Great Gulf - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	22.8	20.2	12.0		23.3	24.8	21.6	21.6	21.5	21.1	21.3	16.8	
Sulfate Bext	76.6	52.5	4.5	80.9	100.1	59.0	66.4	58.7	61.5	59.2	30.8		
Nitrate Bext	3.0	1.8	1.1	3.0	2.9	2.5	3.6	1.5	3.3	1.7	0.9		
POM Bext	14.4	10.5	13.3	11.0	20.1	15.2	11.2	13.0	10.6	11.0	7.3		
EC Bext	3.9	3.2	0.4	3.9	4.4	4.0	3.4	4.0	3.5	3.5	1.9		
Soil Bext	0.6	0.4	0.5	0.6	0.7	0.4	0.5	0.2	0.4	0.5	0.3		
Coarse Mass Bext	3.0	2.6	3.3	3.0	2.7	2.8	3.4	2.2	3.0	2.8	2.2		
Sea Salt Bext	0.1	0.2	0.1	0.0	0.2	0.0	0.3	0.2	0.1	0.3	0.0		




Great Gulf - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.7	6.6	3.7	8.3	7.8	6.9	7.7	6.9	6.4	6.9	6.2		
Sulfate Bext	5.8	4.8	0.7	6.5	6.0	4.8	5.9	5.3	4.7	5.2	4.0		
Nitrate Bext	1.0	0.6	0.4	1.1	1.0	0.7	1.0	0.8	0.4	0.6	0.6		
POM Bext	2.0	1.5	1.6	2.1	2.0	1.9	2.0	1.7	1.6	1.5	1.5		
EC Bext	0.8	0.7	0.1	0.9	0.8	0.8	0.8	0.8	0.7	0.7	0.5		
Soil Bext	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1		
Coarse Mass Bext	0.9	0.7	0.6	1.1	0.7	1.0	0.7	0.6	0.7	0.7	0.8		
Sea Salt Bext	0.2	0.2	0.1	0.0	0.4	0.0	0.2	0.1	0.1	0.2	0.3		




Brigantine - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	29.0	27.3	12.2	29.0	28.4	29.3	29.8	28.6	29.6	28.5	26.9		24.3
Sulfate Bext	127.1	107.4	3.2	122.	116.	122.	145.	129.	148.	119.	100.		61.3
Nitrate Bext	15.7	12.2	1.1	19.2	18.1	13.4	17.8	10.0	12.0	8.2	13.3		15.3
POM Bext	24.2	14.9	12.8	19.2	16.1	48.9	18.8	18.1	16.0	16.9	13.2		13.7
EC Bext	7.0	6.1	0.3	8.1	6.7	7.9	7.0	5.1	6.8	6.4	5.9		5.4
Soil Bext	1.0	0.7	0.8	0.6	1.1	1.1	0.8	1.2	0.6	0.8	0.7		0.7
Coarse Mass Bext	5.4	7.3	2.6	5.5	8.9	4.3	3.5	4.7	6.5	12.9	4.3		5.3
Sea Salt Bext	0.4	1.2	3.6	0.5	0.1	0.1	0.4	0.9	0.9	0.8	1.6		1.3




Brigantine - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.3	13.9	5.5	14.3	13.8	14.8	14.4	14.4	14.6	15.3	12.7		12.8
Sulfate Bext	14.8	13.5	0.9	14.8	14.0	16.3	15.1	13.9	15.8	16.2	11.3		10.9
Nitrate Bext	3.9	3.6	0.4	3.7	3.7	3.6	3.9	4.4	3.9	4.2	3.3		2.8
POM Bext	4.5	3.6	2.5	4.7	4.3	4.3	4.0	5.4	3.3	4.0	3.3		3.8
EC Bext	2.4	1.9	0.1	3.0	2.4	2.1	1.9	2.8	2.2	2.0	1.6		1.7
Soil Bext	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.3	0.2		0.3
Coarse Mass Bext	3.2	3.3	1.0	3.4	3.6	3.9	2.8	2.2	2.9	5.3	2.4		2.5
Sea Salt Bext	1.4	2.5	0.2	0.7	0.2	2.0	2.6	1.3	3.5	2.6	2.0		2.1




Bandelier - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.2	11.8	6.3	14.6	11.1	12.3	12.7	10.5	12.3	11.6	12.8	11.5	11.0
Sulfate Bext	6.9	8.4	0.6	7.6	7.5	7.1	7.1	5.4	10.6	7.7	8.4	7.3	7.9
Nitrate Bext	2.5	2.4	0.8	2.8	2.2	2.6	2.8	2.2	1.4	2.1	4.1	2.3	2.0
POM Bext	14.2	7.6	4.0	38.1	5.4	9.5	11.6	6.5	8.0	6.9	11.8	5.0	6.3
EC Bext	3.1	2.1	0.3	6.8	2.0	2.5	2.5	1.9	2.2	2.1	2.8	1.5	1.9
Soil Bext	1.1	1.2	1.1	0.9	1.1	1.4	1.2	0.9	0.7	1.0	0.5	2.3	1.5
Coarse Mass Bext	2.9	3.2	3.6	2.4	2.6	3.5	3.4	2.7	2.7	3.4	1.9	4.4	3.5
Sea Salt Bext	0.2	0.1	0.2	0.1	0.7	0.3	0.1	0.1	0.1	0.0	0.0	0.1	0.1




Bandelier - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.0	4.2	1.3	5.0	4.9	5.0	4.9	4.9	4.1	4.6	4.5	3.9	3.6
Sulfate Bext	2.4	2.1	0.4	2.4	2.9	2.5	2.0	2.4	2.0	2.5	2.4	1.7	1.9
Nitrate Bext	0.7	0.6	0.3	0.8	0.6	0.8	0.8	0.7	0.6	0.5	0.6	0.7	0.4
POM Bext	2.1	1.8	0.9	2.0	1.7	2.1	2.4	2.4	1.6	1.7	2.0	1.9	1.5
EC Bext	0.9	0.8	0.1	0.9	0.8	0.9	1.0	0.9	0.9	1.0	0.8	0.7	0.6
Soil Bext	0.3	0.2	0.2	0.3	0.4	0.4	0.2	0.3	0.2	0.3	0.2	0.2	0.3
Coarse Mass Bext	0.9	0.8	0.5	1.0	1.0	1.0	1.0	0.7	0.9	1.0	0.8	0.6	0.7
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0




Bosque del Apache - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.8	13.4	6.7			14.6	13.9	12.9	14.3	13.9	14.2	12.8	11.9
Sulfate Bext	7.5	8.9	0.6			8.7	7.3	6.6	11.8	9.1	10.1	8.0	5.5
Nitrate Bext	3.2	4.2	1.0			3.4	2.1	4.1	1.9	4.5	9.4	2.1	3.1
POM Bext	8.7	6.5	3.1			9.6	8.5	8.0	8.0	7.2	7.2	5.2	5.2
EC Bext	2.6	2.8	0.3			2.5	2.5	2.9	3.4	3.2	3.2	1.8	2.6
Soil Bext	1.9	1.6	1.1			2.8	1.7	1.4	1.5	1.7	1.1	2.3	1.4
Coarse Mass Bext	6.7	5.5	3.6			7.6	8.3	4.2	5.3	5.9	4.4	7.0	4.9
Sea Salt Bext	0.2	0.2	0.2			0.1	0.2	0.3	0.2	0.3	0.2	0.1	0.1




Bosque del Apache - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.3	5.8	2.2			6.5	6.8	5.5	5.9	6.3	5.9	5.2	5.5
Sulfate Bext	2.8	2.6	0.3			2.8	2.9	2.6	2.8	2.9	2.9	2.2	2.4
Nitrate Bext	0.8	0.6	0.3			0.8	0.9	0.8	0.6	0.6	0.7	0.6	0.6
POM Bext	2.1	1.8	0.9			2.3	2.4	1.8	1.7	1.8	1.9	1.7	1.7
EC Bext	1.1	0.9	0.1			1.2	1.2	0.9	1.0	1.1	0.9	0.8	0.9
Soil Bext	0.3	0.4	0.2			0.4	0.4	0.3	0.3	0.5	0.3	0.3	0.4
Coarse Mass Bext	1.6	1.4	0.6			1.9	1.9	0.9	1.6	1.8	1.2	1.1	1.5
Sea Salt Bext	0.1	0.1	0.1			0.0	0.2	0.2	0.1	0.2	0.1	0.1	0.0




Gila - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.1	12.5	6.7		12.1	14.2	14.1	12.0	13.9	12.6	13.0	11.1	11.8
Sulfate Bext	6.9	8.0	0.6		8.3	6.4	6.9	5.8	10.1	7.9	8.8	7.0	6.4
Nitrate Bext	0.9	0.8	0.7		0.6	1.0	1.1	0.9	0.6	0.7	0.9	0.8	0.9
POM Bext	16.0	12.5	4.5		10.2	21.2	18.0	14.5	15.7	15.4	16.2	5.7	9.3
EC Bext	3.2	2.9	0.4		2.3	3.5	3.9	2.9	4.6	3.0	3.3	1.2	2.4
Soil Bext	1.5	1.5	1.2		1.4	1.7	1.5	1.2	0.6	1.6	1.4	2.4	1.4
Coarse Mass Bext	2.8	3.6	3.5		2.9	3.4	2.8	2.3	2.0	3.1	4.4	4.5	4.0
Sea Salt Bext	0.1	0.1	0.2		0.0	0.0	0.2	0.1	0.1	0.1	0.1	0.1	0.1




Gila - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.3	2.7	0.5	3.7	3.4	2.9	3.3	3.1	3.3	2.7	1.9	2.7	
Sulfate Bext	1.9	1.7	0.3	2.3	1.8	1.4	1.9	2.0	1.9	1.6	1.3	1.8	
Nitrate Bext	0.3	0.2	0.2	0.4	0.3	0.2	0.4	0.3	0.3	0.2	0.2	0.2	
POM Bext	1.5	1.1	0.6	1.4	1.6	1.6	1.5	1.2	1.2	1.2	0.8	0.9	
EC Bext	0.6	0.4	0.1	0.6	0.8	0.6	0.6	0.5	0.6	0.4	0.3	0.4	
Soil Bext	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.3	0.2	0.2	0.2	
Coarse Mass Bext	0.4	0.5	0.3	0.6	0.4	0.4	0.3	0.4	0.7	0.5	0.4	0.6	
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	




Guadalupe Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.2	15.9	6.7	17.1	16.6	18.1	18.5	15.6	17.5	15.3	15.7	16.8	14.1
Sulfate Bext	16.5	18.6	0.8	16.6	18.6	20.5	12.4	14.4	26.0	16.6	21.5	14.5	14.4
Nitrate Bext	3.8	2.9	0.9	5.4	3.2	2.8	2.6	5.1	3.8	2.7	3.7	2.6	1.5
POM Bext	6.7	5.9	3.1	7.4	5.0	7.8	7.4	6.2	7.4	5.9	5.8	5.6	4.9
EC Bext	1.3	1.5	0.2	2.0	1.0	1.3	1.0	1.3	2.2	1.6	1.5	1.0	1.2
Soil Bext	4.4	2.7	1.3	2.4	4.1	5.1	7.5	2.9	2.1	2.0	2.0	4.8	2.7
Coarse Mass Bext	16.0	9.9	4.4	13.2	12.7	16.0	28.1	10.1	8.3	8.6	6.0	18.8	7.7
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1




Guadalupe Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.9	5.4	1.0	6.3	6.3	6.4	5.9	4.8	6.1	5.9	5.3	5.4	4.6
Sulfate Bext	3.4	3.1	0.3	3.9	3.9	3.6	2.9	2.9	3.7	3.3	3.0	3.1	2.6
Nitrate Bext	1.1	0.9	0.3	0.9	1.3	1.5	1.1	1.0	1.0	0.7	1.1	0.8	0.9
POM Bext	1.6	1.5	0.7	1.9	1.6	1.9	1.5	1.1	1.8	1.9	1.5	1.5	1.0
EC Bext	0.7	0.7	0.1	0.9	0.7	0.7	0.6	0.6	0.9	0.9	0.7	0.6	0.4
Soil Bext	0.4	0.5	0.1	0.5	0.5	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.6
Coarse Mass Bext	2.0	1.7	0.5	1.6	2.0	2.3	2.7	1.5	1.9	2.0	1.4	1.9	1.5
Sea Salt Bext	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1




Salt Creek - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.0	17.5	6.8	17.6	18.0	18.8	17.7	19.2	17.3		17.6	15.7	
Sulfate Bext	16.7	18.9	0.8	18.7	17.9	13.9	16.5	29.3	16.2		16.1	13.7	
Nitrate Bext	11.1	7.4	0.8	12.2	9.5	10.4	12.4	8.4	6.4		9.3	5.4	
POM Bext	7.5	6.4	3.0	6.1	7.3	8.5	8.1	8.4	6.5		5.8	5.0	
EC Bext	2.3	2.3	0.3	2.5	2.2	2.3	2.3	3.3	2.4		1.6	2.1	
Soil Bext	3.3	2.3	1.0	2.1	2.1	4.9	4.2	1.4	2.3		3.6	2.0	
Coarse Mass Bext	11.5	11.4	4.1	10.3	12.4	16.8	6.4	7.5	13.1		13.8	11.0	
Sea Salt Bext	0.2	0.5	0.2	0.3	0.1	0.1	0.3	1.0	0.3		0.3	0.3	




Salt Creek - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.8	7.3	2.1	7.9	7.9	8.5	7.1	7.9	8.0		6.8	6.4	
Sulfate Bext	3.4	3.4	0.2	3.5	3.3	3.7	3.3	4.2	3.6		3.2	2.6	
Nitrate Bext	2.0	1.3	0.3	2.6	2.0	1.8	1.8	2.2	1.3		1.0	0.7	
POM Bext	2.3	2.0	0.9	2.3	2.2	2.1	2.4	2.2	2.3		1.7	1.8	
EC Bext	1.1	0.9	0.1	1.2	1.1	1.1	1.0	1.1	1.2		0.7	0.7	
Soil Bext	0.7	0.5	0.2	0.4	0.5	1.4	0.5	0.4	0.6		0.5	0.5	
Coarse Mass Bext	2.5	2.7	0.6	2.4	3.1	3.4	1.3	2.0	3.5		2.7	2.7	
Sea Salt Bext	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1		0.0	0.0	




San Pedro Parks - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.2	9.9	5.7	9.3	10.6	11.6	9.3	10.8	9.7	10.7	10.0	8.2	
Sulfate Bext	5.8	6.8	0.7	6.5	6.0	5.3	5.3	9.1	6.3	7.4	6.3	4.7	
Nitrate Bext	1.6	1.2	0.8	1.5	1.8	1.6	1.3	1.3	1.3	1.3	1.0	1.3	1.0
POM Bext	7.7	6.3	4.0	4.8	7.8	11.3	6.8	6.3	6.3	7.2	8.6	5.0	4.4
EC Bext	1.6	1.6	0.3	1.1	1.6	2.6	1.1	2.2	1.6	1.6	1.7	1.2	1.1
Soil Bext	1.5	1.3	1.2	1.6	1.7	1.5	1.0	1.1	1.1	0.9	1.0	2.3	1.3
Coarse Mass Bext	2.7	2.5	3.0	2.0	2.6	4.2	2.1	2.1	2.3	2.5	3.2	2.3	
Sea Salt Bext	0.1	0.1	0.1	0.0	0.3	0.1	0.0	0.1	0.0	0.0	0.0	0.1	0.0



San Pedro Parks - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.5	1.0	-0.7	1.9	1.1	1.4	1.5	0.9	1.4	1.0	0.8	1.1	
Sulfate Bext	1.6	1.5	0.3	1.9	1.4	1.4	1.6	1.3	1.9	1.4	1.2	1.5	
Nitrate Bext	0.4	0.4	0.2	0.5	0.4	0.4	0.4	0.4	0.4	0.3	0.4	0.4	0.4
POM Bext	0.7	0.6	0.4	0.9	0.6	0.7	0.7	0.7	0.5	0.6	0.6	0.6	0.6
EC Bext	0.4	0.2	0.1	0.4	0.3	0.4	0.5	0.2	0.3	0.2	0.2	0.2	0.3
Soil Bext	0.2	0.1	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Coarse Mass Bext	0.3	0.3	0.2	0.3	0.3	0.4	0.2	0.3	0.3	0.3	0.3	0.3	0.3
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0




Wheeler Peak - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.4	9.1	6.1			11.1	11.4	8.7	10.1	9.2	8.6		8.6
Sulfate Bext	5.3	6.2	0.8			5.3	6.1	4.5	7.6	6.3	5.7		5.1
Nitrate Bext	1.6	1.4	0.8			2.1	1.8	1.1	1.4	1.6	1.2		1.5
POM Bext	8.4	4.7	4.2			10.1	10.2	4.7	5.7	4.3	5.0		3.9
EC Bext	2.2	1.6	0.4			2.2	2.4	2.0	2.0	1.2	1.4		1.8
Soil Bext	1.7	1.1	1.2			1.6	2.1	1.6	1.3	1.2	0.7		1.3
Coarse Mass Bext	2.8	2.1	3.2			3.3	3.8	1.2	1.9	2.5	1.9		2.3
Sea Salt Bext	0.5	0.1	0.5			0.3	0.1	1.0	0.1	0.0	0.1		0.1




Wheeler Peak - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.2	0.9	-0.6			1.1	1.5	1.0	0.6	1.3	1.2		0.4
Sulfate Bext	1.2	1.3	0.2			1.2	1.1	1.2	1.1	1.5	1.3		1.3
Nitrate Bext	0.3	0.3	0.2			0.4	0.3	0.4	0.2	0.2	0.3		0.2
POM Bext	0.8	0.7	0.5			0.6	1.0	0.8	0.6	0.9	0.8		0.4
EC Bext	0.4	0.3	0.1			0.4	0.4	0.3	0.2	0.4	0.3		0.2
Soil Bext	0.1	0.1	0.1			0.1	0.1	0.1	0.1	0.1	0.1		0.1
Coarse Mass Bext	0.5	0.3	0.4			0.5	0.6	0.4	0.4	0.3	0.5		0.2
Sea Salt Bext	0.1	0.0	0.0			0.1	0.0	0.0	0.0	0.0	0.0		0.0




White Mountain - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.7	13.2	6.8		14.8	13.9	12.4	13.7	12.7	12.4	14.3	12.9	
Sulfate Bext	10.5	11.9	0.7		13.1	10.7	7.8	15.5	10.5	11.4	11.6	10.2	
Nitrate Bext	3.0	1.8	1.0		3.9	2.9	2.3	1.9	1.4	2.2	2.3	1.2	
POM Bext	9.0	5.4	3.7		8.6	7.0	11.3	6.6	5.2	4.8	4.7	5.5	
EC Bext	1.8	1.4	0.3		1.9	1.5	2.1	2.2	1.7	1.3	1.0	1.0	
Soil Bext	1.9	1.8	1.2		2.0	2.1	1.6	1.1	1.8	1.3	2.7	1.9	
Coarse Mass Bext	6.7	7.4	4.3		7.4	8.9	3.8	3.9	7.3	4.7	12.1	9.2	
Sea Salt Bext	0.2	0.3	0.2		0.1	0.2	0.2	0.1	0.3	0.1	0.6	0.4	




White Mountain - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.6	3.3	0.7			4.1	3.5	3.0	3.0	3.4	3.9	3.0	3.2
Sulfate Bext	1.9	1.8	0.2			2.2	1.9	1.6	1.6	1.8	2.0	1.7	1.8
Nitrate Bext	0.5	0.4	0.2			0.5	0.4	0.4	0.3	0.3	0.5	0.3	0.4
POM Bext	1.3	1.1	0.7			1.6	1.2	1.2	0.9	1.1	1.5	1.0	1.1
EC Bext	0.6	0.4	0.1			0.7	0.6	0.5	0.6	0.5	0.5	0.4	0.3
Soil Bext	0.2	0.2	0.1			0.2	0.2	0.1	0.2	0.3	0.2	0.2	0.3
Coarse Mass Bext	0.9	1.0	0.4			1.0	1.1	0.5	0.9	1.0	1.0	0.9	1.0
Sea Salt Bext	0.0	0.0	0.0			0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Great Smoky Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	30.3	28.5	11.2	30.7	30.6	29.9	30.5	29.6	31.6	30.5	30.6	26.7	23.2
Sulfate Bext	172.7	144.7	3.5	167.8	179.9	168.3	187.3	160.3	205.1	174.2	166.3	109.6	68.6
Nitrate Bext	3.4	3.3	1.0	3.5	3.0	2.7	3.2	4.6	1.8	5.0	2.7	1.7	5.4
POM Bext	20.6	18.0	11.4	35.1	18.9	18.8	15.7	14.6	14.5	16.3	28.3	18.1	12.6
EC Bext	5.7	5.1	0.3	8.1	5.0	4.8	5.8	4.7	6.1	5.6	5.9	4.2	3.9
Soil Bext	0.8	0.8	0.9	0.6	0.8	1.0	0.7	1.0	0.6	0.9	0.9	0.9	0.6
Coarse Mass Bext	1.9	3.7	2.9	2.6	2.4	2.0	1.5	1.2	4.9	4.6	4.2	2.6	2.1
Sea Salt Bext	0.2	0.3	0.4	0.0	0.5	0.0	0.0	0.3	0.1	0.3	0.3	0.2	0.3




Great Smoky Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.6	12.4	4.6	14.8	13.4	13.9	12.0	13.7	14.3	12.7	13.2	11.6	10.1
Sulfate Bext	16.4	14.3	0.8	18.7	15.7	18.7	12.8	16.2	17.7	16.2	15.4	12.4	9.6
Nitrate Bext	3.9	3.2	0.6	4.5	3.5	4.0	3.6	4.1	5.0	2.6	4.6	2.5	1.6
POM Bext	4.9	3.7	2.3	6.3	4.8	4.3	4.4	4.7	4.6	3.6	3.8	3.6	3.0
EC Bext	2.2	1.8	0.1	3.0	2.0	2.0	2.0	2.1	2.3	2.0	1.8	1.6	1.2
Soil Bext	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Coarse Mass Bext	1.3	1.4	0.9	1.5	1.5	1.1	1.0	1.5	1.7	1.2	1.1	1.4	1.5
Sea Salt Bext	0.2	0.2	0.1	0.1	0.3	0.2	0.0	0.4	0.2	0.2	0.3	0.2	0.1




Linville Gorge - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	28.8	27.4	11.2		29.4	28.8	28.4	28.5	31.9		29.9	25.1	22.7
Sulfate Bext	146.0	130.9	4.0		152.3	147.2	140.3	144.3	213.9		159.9	89.6	60.1
Nitrate Bext	2.3	2.7	1.1		3.5	2.3	1.7	1.6	2.7		1.6	1.8	4.7
POM Bext	16.1	17.9	11.0		18.5	15.8	16.9	13.1	13.7		23.5	18.3	15.9
EC Bext	5.2	5.0	0.3		5.9	4.5	5.4	4.8	6.2		5.3	4.3	4.4
Soil Bext	0.8	0.6	0.8		0.8	0.8	0.5	0.9	0.4		0.7	0.7	0.5
Coarse Mass Bext	2.2	1.9	2.5		2.8	2.4	2.1	1.5	1.7		2.2	1.9	1.8
Sea Salt Bext	0.1	0.4	0.3		0.0	0.1	0.0	0.4	0.4		0.3	0.5	0.4




Linville Gorge - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.1	11.3	4.1	11.9	12.1	9.7	10.7	12.6		12.4	10.6	9.6	
Sulfate Bext	11.3	11.5	0.6	12.4	13.1	9.2	10.6	13.7		14.4	9.3	8.4	
Nitrate Bext	2.2	2.4	0.5	2.8	2.3	2.4	1.5	3.2		2.5	2.6	1.2	
POM Bext	3.8	4.0	2.0	4.1	4.5	2.8	3.7	4.5		4.2	4.1	3.2	
EC Bext	1.8	1.7	0.1	2.0	2.2	1.3	1.7	2.2		1.8	1.5	1.2	
Soil Bext	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2		0.2	0.2	0.2	
Coarse Mass Bext	0.8	1.0	0.6	1.0	0.9	0.8	0.6	1.0		1.0	1.0	1.0	
Sea Salt Bext	0.1	0.2	0.1	0.1	0.1	0.0	0.3	0.2		0.2	0.1	0.1	




Shining Rock - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	28.5	26.6	11.5		28.9	29.2	29.8	25.9	29.7	29.4	27.9	24.7	21.3
Sulfate Bext	149.0	122.7	4.7		153.6	162.2	175.3	105.1	167.8	160.9	129.6	98.8	56.5
Nitrate Bext	2.7	2.8	1.2		3.0	3.1	2.2	2.4	2.4	2.4	3.7	1.9	3.8
POM Bext	14.6	14.3	12.3		13.7	15.7	16.4	12.7	12.2	17.6	20.0	10.2	11.3
EC Bext	3.6	3.5	0.3		4.0	2.5	4.6	3.5	4.9	4.4	4.2	1.3	2.5
Soil Bext	0.9	0.7	1.0		0.7	1.1	0.8	0.9	0.5	0.9	0.8	0.7	0.7
Coarse Mass Bext	1.3	1.8	2.1		1.3	1.8	0.6	1.3	1.2	2.2	2.3	1.6	1.9
Sea Salt Bext	0.1	0.2	0.2		0.0	0.0	0.0	0.4	0.1	0.3	0.3	0.2	0.2




Shining Rock - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.7	7.3	2.5	8.9	8.4	6.1	7.4	7.5	7.3	8.0	7.8	5.8	
Sulfate Bext	7.0	6.5	0.4	8.3	8.2	4.8	6.8	7.4	6.6	7.0	6.0	5.2	
Nitrate Bext	1.5	1.2	0.4	2.0	1.5	1.0	1.4	1.3	0.9	1.6	1.3	0.8	
POM Bext	2.1	2.3	1.5	2.4	2.3	1.8	1.8	1.9	2.2	2.5	3.7	1.1	
EC Bext	0.9	0.7	0.1	1.0	1.1	0.7	0.8	0.8	0.9	0.9	0.6	0.5	
Soil Bext	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	
Coarse Mass Bext	0.4	0.4	0.3	0.7	0.5	0.2	0.3	0.4	0.4	0.4	0.5	0.4	
Sea Salt Bext	0.1	0.1	0.0	0.2	0.0	0.0	0.1	0.1	0.1	0.2	0.1	0.1	




Swanquarter - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	24.7	24.6	11.5	25.3	25.7	23.4	24.6		26.2	25.6			21.9
Sulfate Bext	83.5	87.2	2.7	86.0	83.8	66.4	97.8		106.6	99.6			55.3
Nitrate Bext	5.8	3.8	1.3	6.2	7.4	5.7	3.9		3.5	4.1			3.8
POM Bext	14.8	12.2	12.1	16.0	23.2	14.8	5.2		12.8	12.4			11.4
EC Bext	3.7	3.4	0.3	4.5	4.8	4.0	1.6		4.0	3.8			2.6
Soil Bext	0.6	0.8	0.7	0.6	0.6	0.5	0.8		0.7	0.6			1.1
Coarse Mass Bext	3.5	3.4	2.1	2.8	3.7	4.3	3.2		3.8	3.1			3.3
Sea Salt Bext	0.2	1.2	1.4	0.0	0.1	0.1	0.6		0.5	0.6			2.4




Swanquarter - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.0	13.0	5.5	13.1	12.8	10.9	11.2			13.8	12.7		12.6
Sulfate Bext	11.3	12.9	0.8	12.5	12.9	9.5	10.2			14.8	12.5		11.6
Nitrate Bext	2.3	2.5	0.4	3.1	2.5	1.5	2.0			2.7	2.6		2.2
POM Bext	2.3	2.9	1.9	3.1	3.1	1.6	1.4			3.5	2.4		2.9
EC Bext	0.8	1.0	0.1	1.1	1.0	0.6	0.5			1.3	0.8		0.9
Soil Bext	0.2	0.2	0.2	0.2	0.2	0.2	0.3			0.3	0.2		0.3
Coarse Mass Bext	3.2	2.6	1.4	3.1	3.0	3.6	3.0			2.8	2.5		2.4
Sea Salt Bext	1.5	3.0	0.3	2.0	1.4	1.3	1.4			2.9	3.2		3.1




Lostwood - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	19.6	19.6	8.0	19.7	20.6	18.8	18.6	20.2	20.5	19.6	19.1	19.7	18.9
Sulfate Bext	21.4	22.9	1.1	20.0	21.5	20.1	18.6	26.8	29.9	20.2	22.9	20.3	21.3
Nitrate Bext	22.9	21.7	1.1	16.0	29.3	23.3	19.4	26.7	19.0	21.4	20.0	21.9	26.3
POM Bext	11.1	10.1	3.8	17.8	9.2	7.6	9.1	11.6	11.0	14.5	8.0	12.2	5.0
EC Bext	2.8	2.5	0.4	4.5	2.8	2.3	2.4	2.2	3.2	2.8	2.1	2.5	2.0
Soil Bext	0.6	0.6	0.9	0.9	0.7	0.5	0.6	0.4	0.5	0.6	0.6	0.7	0.7
Coarse Mass Bext	3.9	4.0	3.7	5.2	4.7	2.6	4.3	2.7	4.1	4.2	4.4	3.9	3.5
Sea Salt Bext	0.3	0.6	0.5	0.0	0.9	0.1	0.0	0.3	0.2	0.2	0.2	2.1	0.1




Lostwood - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	8.2	8.1	2.9	9.1	8.2	7.9	7.9	7.9	7.6	7.8	8.8	8.2	8.4
Sulfate Bext	4.4	4.8	0.4	5.3	4.7	3.8	3.9	4.3	4.7	4.0	5.4	4.4	5.2
Nitrate Bext	1.9	1.5	0.3	2.4	1.6	1.8	1.7	1.8	1.7	1.8	1.5	1.1	1.7
POM Bext	2.3	1.9	0.7	2.9	1.9	2.0	2.5	2.0	1.6	2.0	2.1	2.2	1.8
EC Bext	0.7	0.7	0.1	0.8	0.7	0.7	0.7	0.5	0.7	0.9	0.7	0.7	0.7
Soil Bext	0.3	0.3	0.2	0.3	0.4	0.4	0.3	0.3	0.2	0.3	0.4	0.3	0.4
Coarse Mass Bext	2.3	2.5	0.6	2.2	2.5	2.4	2.2	2.2	1.6	2.1	3.1	3.1	2.5
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.1




Theodore Roosevelt - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.8	17.6	7.8	18.1	18.0	17.0	18.4	17.5	17.6	17.9		17.6	17.2
Sulfate Bext	17.5	18.4	1.0	16.4	18.8	20.8	17.7	14.0	17.7	17.3		16.6	22.0
Nitrate Bext	13.7	12.3	1.0	13.6	17.7	10.7	10.3	16.4	16.1	9.5		11.8	11.9
POM Bext	10.8	10.3	3.9	11.8	6.7	5.9	16.4	13.4	6.3	14.7		14.7	5.4
EC Bext	2.7	2.6	0.3	3.3	2.7	1.9	3.4	2.5	2.8	3.3		2.5	1.9
Soil Bext	0.9	0.8	1.0	0.8	1.0	1.2	1.0	0.5	0.9	1.0		0.8	0.7
Coarse Mass Bext	4.8	4.7	3.7	5.6	4.5	5.6	5.4	3.0	5.1	5.3		4.1	4.4
Sea Salt Bext	0.1	0.6	0.2	0.0	0.0	0.0	0.0	0.3	0.2	0.1		2.0	0.1




Theodore Roosevelt - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	7.8	6.7	3.0	8.2	7.8	7.8	7.5	7.5	6.8	6.5		6.6	7.0
Sulfate Bext	3.8	3.4	0.4	4.9	3.6	3.8	3.5	3.2	3.6	2.5		3.3	4.1
Nitrate Bext	1.5	0.9	0.3	1.6	1.4	1.9	1.6	1.2	1.0	0.9		0.7	1.0
POM Bext	2.0	1.5	0.7	1.9	1.8	2.2	1.8	2.1	1.6	1.4		1.5	1.5
EC Bext	0.9	0.9	0.1	1.2	0.8	0.8	0.9	0.9	1.0	1.1		0.7	0.6
Soil Bext	0.4	0.3	0.2	0.3	0.5	0.4	0.3	0.4	0.3	0.3		0.3	0.3
Coarse Mass Bext	2.2	1.7	0.7	2.1	2.7	1.7	2.1	2.4	1.3	1.9		1.8	1.8
Sea Salt Bext	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1		0.1	0.0




Wichita Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	23.8	23.0	7.5		23.6	23.6	24.2	25.7	21.8	22.8	21.6		
Sulfate Bext	49.1	53.0	1.2		62.2	43.5	41.7	83.9	42.6	45.5	40.1		
Nitrate Bext	23.7	17.0	1.2		10.3	28.5	32.3	10.3	11.2	27.9	18.7		
POM Bext	17.3	12.7	3.2		14.5	14.8	22.5	17.3	13.4	10.9	9.4		
EC Bext	4.5	4.6	0.3		4.2	4.5	4.7	5.9	5.1	4.1	3.0		
Soil Bext	0.8	1.2	1.1		0.8	0.9	0.7	0.7	1.5	1.1	1.3		
Coarse Mass Bext	4.6	5.4	3.1		5.5	4.2	4.3	5.4	7.3	4.4	4.7		
Sea Salt Bext	0.2	0.4	0.2		0.0	0.0	0.4	0.2	0.4	0.6	0.4		




Wichita Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	9.8	9.9	3.0			9.8	10.0	9.6	10.6	9.7	9.3	9.8	
Sulfate Bext	5.1	6.5	0.2			5.4	5.0	5.0	7.5	5.9	5.8	6.7	
Nitrate Bext	2.8	2.2	0.3			3.2	2.7	2.5	2.4	2.0	1.8	2.7	
POM Bext	3.6	3.0	0.8			3.2	3.7	3.9	3.3	2.7	3.2	2.9	
EC Bext	1.3	1.2	0.1			1.2	1.3	1.3	1.5	1.1	1.2	1.0	
Soil Bext	0.3	0.4	0.2			0.3	0.4	0.3	0.4	0.5	0.3	0.4	
Coarse Mass Bext	2.7	2.9	0.9			2.5	3.3	2.2	3.5	3.5	2.3	2.3	
Sea Salt Bext	0.1	0.1	0.1			0.0	0.1	0.2	0.1	0.1	0.2	0.1	




Crater Lake - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.7	13.8	7.6			15.6	13.1	12.5	11.1	17.9	11.5	15.8	12.7
Sulfate Bext	7.3	8.2	1.9			7.3	7.0	7.8	8.1	7.6	8.9	9.1	7.5
Nitrate Bext	2.6	1.7	2.3			4.5	1.3	2.0	1.7	1.9	1.8	1.8	1.4
POM Bext	21.4	22.5	6.2			36.3	15.7	12.3	6.9	50.9	7.8	30.1	17.0
EC Bext	4.3	3.4	0.5			6.7	3.7	2.6	2.4	6.4	2.0	3.6	2.5
Soil Bext	0.8	0.9	0.9			1.0	0.7	0.8	0.6	0.8	0.8	0.7	1.3
Coarse Mass Bext	2.3	1.8	2.5			3.6	1.9	1.4	1.8	1.9	1.7	1.5	2.2
Sea Salt Bext	0.1	0.1	0.2			0.1	0.0	0.0	0.1	0.1	0.2	0.1	0.2




Crater Lake - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.7	1.6	0.1			1.8	1.7	1.6	1.7	1.5	1.4	1.7	1.6
Sulfate Bext	0.9	1.2	0.2			1.0	0.9	1.0	1.3	1.2	1.2	1.2	1.0
Nitrate Bext	0.3	0.2	0.2			0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.2
POM Bext	0.6	0.4	0.2			0.7	0.6	0.5	0.6	0.3	0.3	0.4	0.5
EC Bext	0.7	0.5	0.1			0.7	0.7	0.5	0.4	0.6	0.3	0.5	0.5
Soil Bext	0.1	0.1	0.1			0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.2	0.1			0.2	0.2	0.2	0.1	0.1	0.2	0.3	0.2
Sea Salt Bext	0.2	0.2	0.1			0.3	0.1	0.2	0.2	0.3	0.2	0.2	0.3




Hells Canyon - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.6	18.1	8.3	19.0	18.2			18.5	18.3	19.8	19.4	15.1	
Sulfate Bext	8.4	6.8	1.1	9.9	6.6			8.6	8.8	5.4	7.3	5.7	
Nitrate Bext	28.5	13.5	2.7	32.9	20.0			32.5	29.1	11.2	6.8	6.9	
POM Bext	15.6	31.4	3.7	11.4	20.7			14.7	13.1	43.9	51.0	17.8	
EC Bext	3.1	5.3	0.4	2.9	3.7			2.5	4.3	7.6	6.6	2.5	
Soil Bext	0.7	0.8	0.9	0.7	0.7			0.6	0.4	1.1	1.2	0.7	
Coarse Mass Bext	1.9	2.9	3.4	1.4	2.0			2.5	1.8	3.8	3.6	2.6	
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0			0.1	0.2	0.2	0.1	0.0	




Hells Canyon - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.5	4.8	2.5	5.5	5.8			5.2	4.7	4.7	5.2	4.6	
Sulfate Bext	1.9	1.9	0.4	2.2	1.9			1.7	2.0	1.7	2.2	1.8	
Nitrate Bext	0.8	0.6	0.2	1.0	0.8			0.5	0.5	0.7	0.6	0.7	
POM Bext	2.0	1.5	0.5	1.6	2.3			2.1	1.4	1.5	1.7	1.4	
EC Bext	0.6	0.4	0.1	0.6	0.7			0.5	0.5	0.4	0.5	0.3	
Soil Bext	0.3	0.2	0.1	0.2	0.3			0.2	0.1	0.2	0.2	0.2	
Coarse Mass Bext	0.8	0.5	0.5	0.7	0.9			0.8	0.5	0.4	0.6	0.5	
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0			0.1	0.1	0.2	0.1	0.1	




Kalmiopsis - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.5	16.4	9.4	15.3	15.8	15.5	15.5	19.0	16.9	15.1	15.6	15.7	
Sulfate Bext	10.3	12.1	1.6	11.0	8.4	10.6	11.3	12.1	12.0	10.6	11.2	14.4	
Nitrate Bext	3.2	3.2	1.9	3.4	2.9	2.9	3.5	3.5	4.1	2.8	2.7	2.7	
POM Bext	14.6	20.8	2.9	12.7	19.4	13.8	12.4	47.4	15.7	12.0	17.0	11.8	
EC Bext	2.5	3.5	0.2	2.3	3.5	2.4	1.8	7.3	2.9	3.2	2.4	1.7	
Soil Bext	0.4	0.3	0.4	0.5	0.4	0.3	0.3	0.4	0.4	0.2	0.4	0.4	
Coarse Mass Bext	2.1	2.4	2.2	2.3	1.9	2.2	2.2	3.0	2.5	2.0	2.0	2.3	
Sea Salt Bext	2.9	3.6	4.7	3.2	1.3	3.5	3.8	2.8	5.3	3.2	2.4	4.3	




Kalmiopsis - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.3	6.4	3.7	6.3	5.8	6.2	6.7	6.9	5.8	6.3	6.7	6.1	
Sulfate Bext	1.6	1.8	0.3	1.9	1.6	1.3	1.5	1.7	1.4	1.9	2.7	1.5	
Nitrate Bext	0.4	0.4	0.2	0.5	0.4	0.4	0.3	0.4	0.3	0.4	0.4	0.4	0.3
POM Bext	2.7	2.6	0.9	2.4	2.2	3.0	3.5	3.5	2.2	2.3	2.6	2.5	
EC Bext	0.8	0.8	0.1	0.8	0.6	0.9	1.1	1.1	0.7	0.7	0.6	0.7	
Soil Bext	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1
Coarse Mass Bext	0.6	0.6	0.6	0.7	0.5	0.5	0.5	0.6	0.5	0.6	0.5	0.5	
Sea Salt Bext	0.7	0.8	0.3	0.6	0.7	0.6	0.9	0.7	0.9	0.9	0.9	0.8	




Mount Hood - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.9	13.7	8.4		13.6	13.9	16.4	15.6	12.9	15.2	13.3	14.0	13.2
Sulfate Bext	11.3	10.3	1.9		11.6	10.4	10.7	12.5	9.5	10.1	10.3	11.6	9.9
Nitrate Bext	5.5	4.2	2.3		4.5	5.2	5.8	6.6	5.7	3.8	4.6	3.1	3.8
POM Bext	14.6	12.4	5.6		7.7	9.5	26.7	14.4	7.5	22.9	7.6	15.6	8.6
EC Bext	2.9	2.3	0.5		2.2	2.3	4.3	2.8	2.3	3.2	2.2	2.3	1.8
Soil Bext	0.7	0.6	0.8		1.2	0.6	0.5	0.6	0.3	0.7	0.6	0.7	0.7
Coarse Mass Bext	2.5	2.3	2.7		2.4	2.8	2.7	2.0	1.3	2.6	2.6	2.0	2.9
Sea Salt Bext	0.1	0.7	0.2		0.0	0.0	0.0	0.3	0.6	2.0	0.2	0.1	0.4




Mount Hood - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.2	1.7	0.9	3.1	2.3	1.5	1.7	2.0	1.8	1.7	1.7	1.7	1.5
Sulfate Bext	1.2	1.0	0.3	1.8	1.2	0.7	0.9	1.2	1.0	0.8	0.8	1.0	0.8
Nitrate Bext	0.4	0.3	0.2	0.8	0.4	0.3	0.4	0.3	0.2	0.2	0.3	0.2	0.2
POM Bext	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1
EC Bext	0.2	0.1	0.0	0.3	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1
Soil Bext	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Coarse Mass Bext	0.1	0.1	0.1	0.1	0.2	0.1	0.0	0.1	0.1	0.1	0.2	0.1	0.1
Sea Salt Bext	0.3	0.3	0.1	0.4	0.4	0.2	0.2	0.3	0.4	0.4	0.3	0.2	0.2




Starkey - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.6	16.2	8.9	19.7	18.7	18.5	17.4	19.1	17.1	14.6	15.0	15.3	
Sulfate Bext	7.7	7.8	1.3	8.7	7.0	6.4	8.9	9.0	7.5	7.5	8.1	6.6	
Nitrate Bext	15.8	10.3	2.9	21.1	17.7	5.9	18.4	27.1	6.2	4.1	5.7	8.5	
POM Bext	24.1	19.3	4.0	27.4	24.5	28.3	16.3	20.7	28.7	16.1	15.4	15.5	
EC Bext	4.2	3.6	0.4	4.8	5.2	3.8	3.0	4.8	4.5	3.2	2.6	2.8	
Soil Bext	1.3	0.9	1.2	1.0	0.7	2.3	1.1	0.5	1.5	0.9	0.8	1.0	
Coarse Mass Bext	4.8	3.3	4.3	2.6	3.3	9.4	3.8	1.8	5.0	3.5	2.9	3.5	
Sea Salt Bext	0.1	0.1	0.6	0.0	0.0	0.4	0.2	0.3	0.1	0.1	0.0	0.1	




Starkey - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.5	3.6	1.5	5.4	5.5	3.1	3.9	4.0	3.2	3.6	3.7	3.5	
Sulfate Bext	2.0	1.9	0.4	2.6	2.4	1.2	1.8	2.0	1.7	1.9	1.9	2.0	
Nitrate Bext	0.7	0.6	0.2	0.8	0.8	0.5	0.6	0.4	0.5	0.7	0.5	0.6	
POM Bext	1.7	0.9	0.5	2.1	2.3	1.2	1.4	1.2	0.6	0.8	1.1	0.9	
EC Bext	0.5	0.3	0.1	0.8	0.8	0.3	0.4	0.4	0.3	0.3	0.3	0.2	
Soil Bext	0.1	0.1	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
Coarse Mass Bext	0.5	0.4	0.3	0.7	0.8	0.3	0.4	0.6	0.3	0.3	0.4	0.3	
Sea Salt Bext	0.2	0.3	0.1	0.3	0.2	0.1	0.3	0.2	0.4	0.4	0.2	0.3	




Three Sisters - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.3	16.2	8.8	14.9	14.1	15.7	16.5	15.5	14.5	17.5	15.1	15.8	18.1
Sulfate Bext	11.8	10.8	2.0	13.4	12.8	11.4	10.7	10.6	10.0	11.5	10.9	12.4	9.3
Nitrate Bext	2.7	2.2	2.3	3.2	2.1	2.2	3.6	2.4	2.6	2.7	1.9	1.6	2.0
POM Bext	17.1	17.1	5.0	11.7	9.3	18.7	23.0	22.8	11.7	23.9	15.7	20.4	14.0
EC Bext	3.2	3.3	0.4	3.2	2.4	3.7	3.6	3.1	3.5	4.5	2.9	3.2	2.6
Soil Bext	0.6	1.4	0.6	0.4	0.9	0.7	0.6	0.5	0.7	1.1	0.7	0.7	3.9
Coarse Mass Bext	2.7	7.6	2.8	2.2	2.3	2.8	3.5	2.8	3.8	6.3	4.8	2.8	20.3
Sea Salt Bext	0.2	0.4	0.5	0.0	0.5	0.0	0.3	0.2	0.1	0.9	0.2	0.2	0.5




Three Sisters - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.0	3.0	1.9	3.7	3.8	2.8	2.3	2.6	2.9	2.8	2.9	3.3	3.2
Sulfate Bext	1.1	1.2	0.2	1.3	1.7	1.1	0.6	0.9	1.2	1.0	1.0	1.4	1.4
Nitrate Bext	0.3	0.2	0.2	0.4	0.6	0.2	0.3	0.2	0.2	0.2	0.3	0.2	0.2
POM Bext	0.5	0.4	0.2	0.5	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.6	0.5
EC Bext	0.2	0.2	0.0	0.4	0.2	0.2	0.1	0.1	0.2	0.2	0.1	0.2	0.2
Soil Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Coarse Mass Bext	0.2	0.2	0.2	0.5	0.3	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.1
Sea Salt Bext	0.2	0.3	0.1	0.4	0.3	0.2	0.0	0.3	0.3	0.3	0.3	0.3	0.3




Cape Romain - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	26.5	26.5	12.1	28.8	25.7	25.4	24.7	27.8	29.9	27.9	26.9	24.5	23.0
Sulfate Bext	96.5	107.6	2.4	123.4	83.3	86.9	76.4	112.7	165.8	122.9	110.8	77.5	61.0
Nitrate Bext	5.3	4.2	1.2	6.8	6.4	5.2	4.2	4.0	3.3	3.9	4.4	4.3	5.0
POM Bext	23.8	15.9	11.1	27.6	23.1	20.2	17.3	30.7	17.4	17.5	17.9	14.0	12.5
EC Bext	5.7	4.7	0.3	8.1	5.2	5.5	4.8	5.1	5.3	5.9	4.9	4.0	3.2
Soil Bext	0.6	0.8	0.7	0.5	0.5	0.7	0.4	0.8	0.6	0.9	0.8	0.8	0.7
Coarse Mass Bext	3.2	3.5	2.6	3.7	2.6	2.6	4.9	2.4	2.9	3.2	3.9	3.6	3.9
Sea Salt Bext	0.2	1.3	4.0	0.2	0.2	0.0	0.0	0.4	0.5	0.3	1.0	2.2	2.7




Cape Romain - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.3	15.0	5.9	15.5	13.8	13.6	13.7	14.9	16.3	15.5	14.8	15.0	13.3
Sulfate Bext	15.7	18.0	1.2	17.1	15.4	14.2	14.3	17.3	21.3	20.4	17.3	17.3	13.4
Nitrate Bext	3.1	3.1	0.6	4.5	3.1	2.7	2.4	3.0	3.8	2.8	2.6	3.8	2.4
POM Bext	4.7	4.8	2.6	5.5	4.2	3.4	4.9	5.2	5.0	4.4	5.2	4.8	4.3
EC Bext	1.7	1.8	0.1	2.0	1.8	1.1	1.6	1.8	2.2	2.2	1.7	1.5	1.4
Soil Bext	0.3	0.4	0.2	0.4	0.3	0.2	0.2	0.3	0.6	0.5	0.4	0.3	0.3
Coarse Mass Bext	2.9	2.6	1.2	3.5	2.4	2.7	3.4	2.5	3.0	2.6	2.6	2.4	2.3
Sea Salt Bext	2.2	2.9	0.3	2.7	1.2	3.3	1.0	2.5	3.6	2.9	2.7	2.9	2.2




Badlands - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.1	16.6	8.1	18.1	17.6	16.2	17.8	15.9	16.3	17.5	16.6	17.3	15.1
Sulfate Bext	18.9	18.8	1.2	19.0	23.3	17.4	16.2	18.4	21.8	18.2	17.8	17.2	19.1
Nitrate Bext	5.8	5.4	0.9	3.6	7.4	4.7	7.2	6.3	6.1	5.3	6.2	5.3	4.0
POM Bext	11.8	10.5	4.1	18.7	7.9	7.4	16.2	8.6	5.8	13.2	10.6	17.2	5.7
EC Bext	2.6	1.9	0.3	4.0	2.6	1.9	2.9	1.6	2.1	2.7	2.0	1.6	1.2
Soil Bext	1.0	1.0	1.0	1.0	1.3	1.1	0.7	0.7	0.8	1.2	0.9	1.2	0.7
Coarse Mass Bext	5.9	5.1	4.0	6.1	5.9	7.3	7.0	3.5	3.9	6.8	4.7	6.3	4.0
Sea Salt Bext	0.2	0.5	0.2	0.0	0.0	0.5	0.3	0.1	0.1	0.1	0.2	2.2	0.2




Badlands - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.9	6.8	2.9	7.4	7.5	6.7	6.3	6.6	7.0	6.2	7.1	7.6	6.1
Sulfate Bext	3.4	3.5	0.4	4.3	3.9	2.6	3.0	3.0	4.0	3.0	3.6	3.5	3.2
Nitrate Bext	1.2	1.1	0.4	1.4	1.6	0.9	0.8	1.3	1.2	0.8	1.5	1.1	1.0
POM Bext	1.8	2.2	0.7	1.9	1.9	1.8	1.8	1.6	2.0	1.4	1.8	4.1	1.5
EC Bext	0.8	0.6	0.1	0.8	0.9	0.9	0.6	0.7	0.8	0.6	0.6	0.3	0.5
Soil Bext	0.3	0.3	0.2	0.3	0.4	0.4	0.2	0.3	0.3	0.4	0.4	0.3	0.3
Coarse Mass Bext	1.5	1.2	0.6	1.3	1.6	1.9	1.3	1.4	1.1	1.4	1.5	1.1	1.0
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0




Wind Cave - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.8	15.5	7.7	16.1	15.3	16.6	16.1	15.1	15.7	16.3	16.5	15.1	13.9
Sulfate Bext	13.2	13.7	1.1	11.6	14.3	14.7	12.7	12.8	14.6	16.2	14.2	10.1	13.4
Nitrate Bext	7.0	5.9	1.2	4.7	9.3	5.9	6.2	8.9	8.7	5.6	6.9	4.5	3.7
POM Bext	13.2	13.4	4.4	17.0	7.9	16.6	15.1	9.5	11.2	14.0	16.1	17.1	8.4
EC Bext	2.9	2.6	0.4	3.7	2.2	3.1	2.4	3.2	2.8	3.3	2.8	2.5	1.6
Soil Bext	0.9	1.0	1.0	1.0	0.8	1.0	0.7	0.8	0.8	1.0	1.0	1.0	1.2
Coarse Mass Bext	3.5	3.8	3.8	3.9	3.5	4.1	4.0	2.1	3.1	4.3	3.6	4.4	3.4
Sea Salt Bext	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.0	1.0	0.2




Wind Cave - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.1	4.6	1.9	5.6	5.1	5.2	5.0	4.8	4.7	4.5	5.5	4.3	3.8
Sulfate Bext	2.6	2.5	0.3	3.1	2.6	2.3	2.5	2.4	2.7	2.3	3.2	2.1	2.1
Nitrate Bext	0.8	0.6	0.2	1.1	0.9	0.7	0.5	0.8	0.7	0.5	0.8	0.7	0.6
POM Bext	1.4	1.2	0.6	1.7	1.3	1.4	1.5	1.3	1.1	1.1	1.6	1.3	0.9
EC Bext	0.7	0.5	0.1	0.9	0.6	0.7	0.6	0.7	0.7	0.6	0.6	0.4	0.4
Soil Bext	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Coarse Mass Bext	1.1	0.8	0.6	0.7	1.0	1.5	1.1	0.9	0.8	1.1	0.9	0.8	0.6
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




Great Smoky Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	30.3	28.5	11.2	30.7	30.6	29.9	30.5	29.6	31.6	30.5	30.6	26.7	23.2
Sulfate Bext	172.7	144.7	3.5	167.8	179.9	168.3	187.3	160.3	205.1	174.2	166.3	109.6	68.6
Nitrate Bext	3.4	3.3	1.0	3.5	3.0	2.7	3.2	4.6	1.8	5.0	2.7	1.7	5.4
POM Bext	20.6	18.0	11.4	35.1	18.9	18.8	15.7	14.6	14.5	16.3	28.3	18.1	12.6
EC Bext	5.7	5.1	0.3	8.1	5.0	4.8	5.8	4.7	6.1	5.6	5.9	4.2	3.9
Soil Bext	0.8	0.8	0.9	0.6	0.8	1.0	0.7	1.0	0.6	0.9	0.9	0.9	0.6
Coarse Mass Bext	1.9	3.7	2.9	2.6	2.4	2.0	1.5	1.2	4.9	4.6	4.2	2.6	2.1
Sea Salt Bext	0.2	0.3	0.4	0.0	0.5	0.0	0.0	0.3	0.1	0.3	0.3	0.2	0.3




Great Smoky Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	13.6	12.4	4.6	14.8	13.4	13.9	12.0	13.7	14.3	12.7	13.2	11.6	10.1
Sulfate Bext	16.4	14.3	0.8	18.7	15.7	18.7	12.8	16.2	17.7	16.2	15.4	12.4	9.6
Nitrate Bext	3.9	3.2	0.6	4.5	3.5	4.0	3.6	4.1	5.0	2.6	4.6	2.5	1.6
POM Bext	4.9	3.7	2.3	6.3	4.8	4.3	4.4	4.7	4.6	3.6	3.8	3.6	3.0
EC Bext	2.2	1.8	0.1	3.0	2.0	2.0	2.0	2.1	2.3	2.0	1.8	1.6	1.2
Soil Bext	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Coarse Mass Bext	1.3	1.4	0.9	1.5	1.5	1.1	1.0	1.5	1.7	1.2	1.1	1.4	1.5
Sea Salt Bext	0.2	0.2	0.1	0.1	0.3	0.2	0.0	0.4	0.2	0.2	0.3	0.2	0.1




Big Bend - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.3	16.7	7.2	17.3	18.2	17.2	16.5	18.2	16.9	17.4	16.1	15.0	
Sulfate Bext	26.1	27.8	0.6	25.5	30.2	21.9	26.8	34.4	26.5	35.6	23.4	19.1	
Nitrate Bext	2.0	1.6	0.8	1.6	1.6	3.6	1.4	1.3	2.2	1.6	1.4	1.8	
POM Bext	8.0	6.1	3.2	5.6	10.1	10.2	6.1	7.6	6.3	5.7	5.9	5.1	
EC Bext	2.1	2.2	0.3	2.1	2.3	2.3	1.8	2.9	2.6	2.1	1.7	1.5	
Soil Bext	2.5	1.8	1.2	2.8	2.7	2.5	2.2	2.0	1.2	1.5	2.3	2.1	
Coarse Mass Bext	7.0	5.3	4.2	9.5	7.7	6.5	4.4	4.9	5.7	3.9	6.3	5.5	
Sea Salt Bext	0.1	0.1	0.4	0.2	0.0	0.0	0.2	0.1	0.1	0.1	0.1	0.0	




Big Bend - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.8	5.9	1.6	7.1	5.7	5.7	4.6	6.3	6.3	5.7	5.5	5.7	
Sulfate Bext	3.5	3.2	0.2	5.0	3.5	3.0	2.6	3.7	3.5	3.3	2.8	2.8	
Nitrate Bext	0.5	0.4	0.2	0.7	0.6	0.4	0.4	0.5	0.5	0.4	0.3	0.4	
POM Bext	1.5	1.5	0.7	1.9	1.2	1.4	1.3	1.6	1.5	1.5	1.4	1.5	
EC Bext	0.7	0.6	0.1	0.9	0.6	0.6	0.5	0.7	0.8	0.7	0.5	0.5	
Soil Bext	0.4	0.5	0.1	0.4	0.5	0.5	0.2	0.5	0.5	0.3	0.4	0.6	
Coarse Mass Bext	1.4	1.9	0.5	1.6	1.4	1.6	0.9	1.9	2.1	1.5	2.0	1.9	
Sea Salt Bext	0.1	0.0	0.0	0.0	0.0	0.4	0.1	0.1	0.0	0.1	0.0	0.0	




Guadalupe Mountains - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.2	15.9	6.7	17.1	16.6	18.1	18.5	15.6	17.5	15.3	15.7	16.8	14.1
Sulfate Bext	16.5	18.6	0.8	16.6	18.6	20.5	12.4	14.4	26.0	16.6	21.5	14.5	14.4
Nitrate Bext	3.8	2.9	0.9	5.4	3.2	2.8	2.6	5.1	3.8	2.7	3.7	2.6	1.5
POM Bext	6.7	5.9	3.1	7.4	5.0	7.8	7.4	6.2	7.4	5.9	5.8	5.6	4.9
EC Bext	1.3	1.5	0.2	2.0	1.0	1.3	1.0	1.3	2.2	1.6	1.5	1.0	1.2
Soil Bext	4.4	2.7	1.3	2.4	4.1	5.1	7.5	2.9	2.1	2.0	2.0	4.8	2.7
Coarse Mass Bext	16.0	9.9	4.4	13.2	12.7	16.0	28.1	10.1	8.3	8.6	6.0	18.8	7.7
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.1




Guadalupe Mountains - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.9	5.4	1.0	6.3	6.3	6.4	5.9	4.8	6.1	5.9	5.3	5.4	4.6
Sulfate Bext	3.4	3.1	0.3	3.9	3.9	3.6	2.9	2.9	3.7	3.3	3.0	3.1	2.6
Nitrate Bext	1.1	0.9	0.3	0.9	1.3	1.5	1.1	1.0	1.0	0.7	1.1	0.8	0.9
POM Bext	1.6	1.5	0.7	1.9	1.6	1.9	1.5	1.1	1.8	1.9	1.5	1.5	1.0
EC Bext	0.7	0.7	0.1	0.9	0.7	0.7	0.6	0.6	0.9	0.9	0.7	0.6	0.4
Soil Bext	0.4	0.5	0.1	0.5	0.5	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.6
Coarse Mass Bext	2.0	1.7	0.5	1.6	2.0	2.3	2.7	1.5	1.9	2.0	1.4	1.9	1.5
Sea Salt Bext	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1




Bryce Canyon - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.6	11.9	6.8	10.8	11.3	13.2	11.1	11.8	11.8	10.7	12.7	11.0	13.3
Sulfate Bext	5.2	5.0	0.6	6.5	6.3	4.2	4.4	4.8	6.2	5.1	4.9	5.0	4.0
Nitrate Bext	2.5	2.2	0.7	2.6	3.8	1.4	1.4	3.4	2.3	2.3	3.8	1.5	1.4
POM Bext	9.4	11.8	4.3	6.1	7.5	14.1	11.1	8.0	10.5	7.5	13.0	7.5	20.7
EC Bext	2.4	2.5	0.4	2.1	1.9	2.3	2.4	3.2	2.6	2.1	3.0	1.5	3.5
Soil Bext	1.2	1.3	1.2	1.0	1.4	1.5	1.1	1.2	0.8	0.8	1.2	2.4	1.3
Coarse Mass Bext	4.0	3.1	4.1	2.6	3.1	6.1	4.3	3.9	2.5	2.6	2.9	3.7	3.7
Sea Salt Bext	0.1	0.1	0.1	0.0	0.0	0.3	0.0	0.1	0.0	0.0	0.1	0.2	0.1




Bryce Canyon - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.8	2.1	0.6	3.0	3.1	2.6	2.4	2.8	1.9	2.4	2.4	1.7	2.1
Sulfate Bext	1.5	1.4	0.2	1.6	1.8	1.4	1.1	1.5	1.2	1.7	1.3	1.1	1.4
Nitrate Bext	0.7	0.5	0.3	0.7	0.8	0.8	0.5	0.7	0.4	0.4	0.7	0.4	0.6
POM Bext	1.0	0.7	0.5	1.2	0.8	1.0	1.1	1.1	0.8	0.7	0.9	0.6	0.6
EC Bext	0.4	0.2	0.1	0.4	0.4	0.4	0.3	0.5	0.3	0.3	0.2	0.2	0.2
Soil Bext	0.1	0.1	0.1	0.2	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.2
Coarse Mass Bext	0.5	0.4	0.4	0.5	0.7	0.4	0.5	0.3	0.3	0.4	0.4	0.4	0.4
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0




Canyonlands - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.2	11.0	6.4	11.2	10.9	12.7	11.8	9.6	10.6	10.5	11.4	11.1	11.5
Sulfate Bext	5.6	5.3	0.6	6.3	6.8	4.0	5.5	5.3	5.8	5.6	5.0	5.2	4.8
Nitrate Bext	3.0	3.3	0.8	3.4	3.9	2.1	2.1	3.2	3.9	3.4	3.7	1.6	3.7
POM Bext	7.1	6.2	3.9	6.4	4.2	11.0	9.5	4.2	4.7	4.3	8.7	4.4	8.6
EC Bext	1.7	1.6	0.3	2.1	1.4	2.0	1.9	1.3	1.8	1.7	2.1	0.9	1.7
Soil Bext	1.5	1.5	1.0	0.8	1.8	2.8	1.0	0.9	0.9	1.4	0.9	2.6	1.8
Coarse Mass Bext	3.8	4.6	3.6	3.4	3.1	6.0	4.1	2.4	3.0	3.5	3.3	7.8	5.4
Sea Salt Bext	0.1	0.1	0.1	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.1




Canyonlands - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.7	2.8	1.0	3.6	4.1	4.2	3.3	3.5	2.3	2.7	3.0	2.7	3.3
Sulfate Bext	2.2	1.8	0.4	2.1	2.3	2.4	1.9	2.1	1.8	1.9	1.7	1.7	2.0
Nitrate Bext	0.6	0.5	0.3	0.4	0.6	0.8	0.6	0.5	0.4	0.5	0.6	0.6	0.6
POM Bext	1.1	0.7	0.7	1.2	1.2	1.2	1.0	1.1	0.5	0.5	0.7	0.8	0.7
EC Bext	0.4	0.3	0.1	0.4	0.5	0.5	0.4	0.4	0.2	0.4	0.3	0.3	0.3
Soil Bext	0.3	0.2	0.1	0.2	0.4	0.4	0.2	0.3	0.1	0.2	0.3	0.2	0.3
Coarse Mass Bext	1.0	0.7	0.5	1.0	1.1	1.1	0.9	0.7	0.6	0.6	0.9	0.6	0.9
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0




Capitol Reef - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.9	11.3	6.0	11.6	10.8	12.1	9.4	10.6	11.8		11.7	11.4	10.3
Sulfate Bext	5.9	5.7	0.6	6.3	7.1	4.2	5.1	6.7	7.3		5.6	5.1	4.8
Nitrate Bext	3.4	2.7	0.5	3.1	4.7	3.5	2.0	3.8	3.3		3.2	2.2	2.2
POM Bext	5.8	7.6	3.4	7.2	3.8	9.0	4.8	4.4	7.4		9.5	6.9	6.6
EC Bext	1.6	1.8	0.3	2.0	1.4	2.0	1.3	1.4	2.5		2.2	1.3	1.1
Soil Bext	1.3	1.6	1.1	1.0	1.4	1.5	0.8	1.5	0.8		1.3	2.4	1.9
Coarse Mass Bext	3.4	4.1	3.6	4.6	2.5	4.9	2.7	2.1	3.0		3.8	5.1	4.3
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0		0.1	0.2	0.1




Capitol Reef - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	4.1	2.7	1.3	4.5	4.5	4.2	3.1	4.1	2.6		3.1	2.5	2.7
Sulfate Bext	1.9	1.6	0.3	1.8	2.1	1.8	1.4	2.2	1.6		1.6	1.4	1.9
Nitrate Bext	1.0	0.6	0.5	1.1	1.1	1.1	0.7	0.9	0.4		0.6	0.6	0.6
POM Bext	1.4	0.9	0.7	1.7	1.1	1.6	1.0	1.4	1.0		1.1	0.9	0.5
EC Bext	0.6	0.3	0.1	0.6	0.6	0.8	0.5	0.5	0.4		0.4	0.3	0.2
Soil Bext	0.3	0.2	0.2	0.2	0.4	0.2	0.4	0.4	0.1		0.3	0.1	0.3
Coarse Mass Bext	1.0	0.6	0.5	1.2	1.4	0.8	0.7	0.7	0.4		0.7	0.6	0.7
Sea Salt Bext	0.1	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.0		0.1	0.1	0.0




Lye Brook - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	24.4	23.0	11.7	23.5	26.3	25.5	24.0	22.9	26.0	22.3	25.3		18.4
Sulfate Bext	87.3	79.0	4.8	68.7	117.4	88.2	87.7	74.4	117.9	62.4	97.1		38.4
Nitrate Bext	9.1	5.6	1.4	10.7	7.1	11.3	7.9	8.4	2.0	8.9	4.5		6.9
POM Bext	15.3	10.8	14.8	10.0	12.1	31.2	14.0	9.4	11.8	10.2	14.7		6.5
EC Bext	4.8	4.0	0.4	5.2	4.7	5.7	4.8	3.4	4.4	4.2	4.9		2.5
Soil Bext	0.6	0.5	0.7	0.4	0.8	0.8	0.5	0.7	0.4	0.4	0.7		0.5
Coarse Mass Bext	1.8	2.5	2.0	1.6	2.1	1.4	2.2	1.7	2.1	2.3	3.1		2.2
Sea Salt Bext	0.1	0.3	0.1	0.0	0.0	0.1	0.2	0.2	0.2	0.2	0.2		0.5




Lye Brook - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.4	5.2	2.8	6.5	6.5	6.4	5.8	6.6	5.7	5.2	5.7		4.1
Sulfate Bext	4.4	3.4	0.4	5.1	4.4	4.7	3.7	4.4	4.0	3.4	4.1		2.2
Nitrate Bext	1.2	0.8	0.3	1.2	1.2	1.2	0.9	1.3	0.9	0.8	0.6		0.8
POM Bext	1.3	0.8	1.0	1.1	1.2	1.2	1.2	1.7	1.0	0.8	0.9		0.4
EC Bext	0.6	0.4	0.1	0.6	0.7	0.6	0.6	0.6	0.5	0.5	0.5		0.3
Soil Bext	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.1		0.1
Coarse Mass Bext	0.5	0.5	0.4	0.4	0.6	0.5	0.6	0.5	0.6	0.6	0.5		0.4
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.1	0.1		0.1




James River Face - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	29.1	27.3	11.1	29.5	30.4	28.4	28.2	30.5	29.0	28.5	25.5	22.9	
Sulfate Bext	136.4	116.7	3.2	133.4	161.8	128.1	122.1	176.1	135.8	132.0	78.6	60.8	
Nitrate Bext	7.8	4.6	1.1	12.2	4.3	7.4	7.2	3.8	4.3	5.1	5.2	4.6	
POM Bext	22.8	20.0	11.5	24.8	25.8	20.0	20.8	15.0	23.0	19.9	26.9	15.4	
EC Bext	7.4	7.5	0.3	8.5	7.0	7.2	6.9	8.6	8.7	7.3	7.1	5.8	
Soil Bext	0.8	0.7	0.8	0.9	1.1	0.6	0.8	0.5	0.6	0.8	0.7	0.6	
Coarse Mass Bext	3.0	3.0	2.8	3.5	3.0	2.8	2.8	3.7	2.6	3.0	2.8	2.7	
Sea Salt Bext	0.2	0.3	0.2	0.5	0.0	0.0	0.5	0.2	0.4	0.3	0.3	0.1	




James River Face - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	14.2	13.6	4.4	14.5	15.6	12.8	13.8	14.9	14.8	13.8	13.0	11.6	
Sulfate Bext	17.5	16.5	0.8	19.2	21.2	14.1	15.5	20.0	19.3	17.3	14.9	11.0	
Nitrate Bext	3.7	3.3	0.5	3.3	4.9	3.8	2.9	4.3	3.5	3.2	3.1	2.3	
POM Bext	5.6	4.7	2.1	5.4	6.2	4.6	6.0	5.3	5.6	4.3	4.7	3.8	
EC Bext	3.0	2.8	0.1	2.8	3.4	2.5	3.2	3.2	3.4	2.7	2.7	2.1	
Soil Bext	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.2
Coarse Mass Bext	1.2	1.5	0.8	1.2	1.3	1.2	1.2	1.4	1.8	1.5	1.3	1.6	
Sea Salt Bext	0.1	0.1	0.1	0.1	0.0	0.0	0.2	0.1	0.2	0.2	0.1	0.2	




Shenandoah - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	29.3	27.3	11.4	28.5	29.2	30.5	28.9	29.3	30.8	29.3	28.8	25.6	21.8
Sulfate Bext	155.5	131.2	3.6	135.9	156.2	174.8	153.0	157.6	193.4	158.0	146.6	95.8	62.0
Nitrate Bext	5.8	3.5	0.6	4.5	6.2	5.9	6.3	6.1	4.5	2.7	3.3	3.5	3.6
POM Bext	16.1	13.4	13.7	16.7	15.1	20.4	14.7	13.7	10.8	13.9	16.9	17.1	8.5
EC Bext	5.7	4.7	0.3	6.5	5.9	5.5	5.6	5.1	5.3	5.2	5.2	4.4	3.5
Soil Bext	0.7	0.7	0.8	0.4	0.6	1.1	0.5	1.0	0.5	0.7	0.8	0.8	0.6
Coarse Mass Bext	2.5	2.4	3.0	2.8	2.8	3.3	2.0	1.8	2.1	2.5	2.7	2.5	2.3
Sea Salt Bext	0.1	0.3	0.2	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.4	0.3	0.2




Shenandoah - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	10.9	9.7	3.1	11.1	13.2	11.5	9.5	9.4	10.5	10.6	11.1	8.2	8.2
Sulfate Bext	11.2	10.1	0.6	8.7	15.8	11.9	10.0	9.4	11.6	10.8	12.2	8.1	7.7
Nitrate Bext	4.2	2.9	0.6	4.7	5.4	4.9	2.9	3.0	2.9	3.2	4.3	2.3	1.8
POM Bext	2.9	2.2	1.6	4.0	3.1	2.8	2.3	2.1	2.8	2.4	2.2	1.5	2.0
EC Bext	1.6	1.3	0.1	2.1	1.8	1.6	1.3	1.2	1.7	1.5	1.4	0.9	1.0
Soil Bext	0.2	0.1	0.1	0.2	0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1
Coarse Mass Bext	1.1	0.9	0.7	1.4	1.4	0.9	1.1	0.7	0.8	1.1	0.9	0.8	1.1
Sea Salt Bext	0.1	0.2	0.1	0.1	0.3	0.0	0.1	0.3	0.1	0.3	0.2	0.1	0.1




Virgin Islands - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.0	18.4	10.7	16.8	16.4	17.4	17.6	17.9	18.6			18.2	18.8
Sulfate Bext	10.3	13.9	1.5	9.8	9.7	12.4	9.3	14.5	14.2			11.6	15.3
Nitrate Bext	2.1	2.8	0.9	2.3	1.0	2.5	2.6	2.9	2.7			2.9	2.7
POM Bext	1.8	1.4	1.7	2.1	3.0	1.3	0.9	0.9	0.9			0.8	2.8
EC Bext	1.5	1.0	0.3	1.4	2.1	1.3	1.4	0.9	0.6			0.3	2.3
Soil Bext	5.9	8.2	1.4	5.8	7.0	4.1	6.7	7.8	9.3			7.4	8.4
Coarse Mass Bext	15.7	15.0	3.3	15.1	14.9	15.2	17.6	14.8	17.2			14.1	13.9
Sea Salt Bext	7.5	11.2	9.4	6.5	3.8	9.7	9.9	8.4	9.9			15.7	10.7




Virgin Islands - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	8.5	9.5	4.4		9.1	8.3	8.1	8.6	9.2	9.9		8.9	9.9
Sulfate Bext	4.1	5.1	1.0		4.2	4.1	4.2	3.9	4.6	5.4		4.6	5.8
Nitrate Bext	0.7	1.2	0.4		0.7	0.4	0.7	1.2	1.2	1.2		1.1	1.3
POM Bext	0.6	0.5	0.6		1.0	0.5	0.5	0.5	0.2	0.6		0.6	0.5
EC Bext	0.5	0.4	0.1		0.4	0.5	0.5	0.6	0.4	0.5		0.4	0.4
Soil Bext	0.3	0.3	0.1		0.4	0.4	0.2	0.2	0.3	0.2		0.3	0.3
Coarse Mass Bext	4.0	3.2	0.9		4.3	4.4	3.6	3.5	3.2	3.4		2.9	3.3
Sea Salt Bext	2.3	4.2	1.4		2.7	1.7	1.9	2.8	4.3	4.6		3.5	4.5




Snoqualmie Pass - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	17.8	16.1	8.4	17.5	17.9	18.7	17.2	16.1	16.7	16.2	14.9	16.6	
Sulfate Bext	17.1	15.0	2.0	19.3	15.8	17.4	15.8	14.3	17.3	14.2	13.7	15.3	
Nitrate Bext	11.6	8.9	2.6	10.1	15.8	10.1	10.2	8.4	7.5	13.8	4.7	10.1	
POM Bext	15.4	10.8	4.4	12.0	15.4	19.8	14.5	11.6	11.4	8.4	10.8	11.9	
EC Bext	4.2	3.3	0.4	4.3	4.4	4.7	3.5	3.4	3.9	3.4	2.9	2.9	
Soil Bext	0.4	0.4	0.5	0.4	0.5	0.4	0.4	0.3	0.5	0.3	0.5	0.4	
Coarse Mass Bext	1.5	1.5	1.7	1.2	1.6	1.9	1.3	1.4	2.1	1.0	1.7	1.4	
Sea Salt Bext	0.4	0.4	1.0	0.0	0.1	1.3	0.3	0.9	0.3	0.3	0.2	0.5	




Snoqualmie Pass - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.5	4.9	2.3		6.5	5.4	4.7	5.4	5.4	4.9	4.9	4.5	4.9
Sulfate Bext	2.6	2.1	0.4		3.6	2.5	1.9	2.3	2.6	1.9	1.9	2.2	2.0
Nitrate Bext	1.2	1.0	0.3		1.5	1.2	0.9	1.0	1.3	1.2	1.0	0.5	0.9
POM Bext	1.1	0.8	0.4		1.3	0.9	0.9	1.1	0.8	0.9	0.7	1.0	0.8
EC Bext	0.9	0.7	0.1		1.1	0.7	0.8	1.0	0.7	0.7	0.8	0.6	0.5
Soil Bext	0.1	0.0	0.1		0.1	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.0
Coarse Mass Bext	0.2	0.1	0.2		0.3	0.2	0.2	0.2	0.1	0.2	0.1	0.1	0.1
Sea Salt Bext	0.5	0.7	0.2		0.4	0.6	0.5	0.6	0.7	0.6	0.8	0.3	1.0




White Pass - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.8	12.7	8.4	12.3	11.9	14.1	12.6	11.6	14.0	10.9	13.0	13.8	
Sulfate Bext	9.9	8.9	2.0	11.7	8.8	9.7	9.5	8.4	9.5	8.1	10.1	8.4	
Nitrate Bext	3.1	2.9	2.4	2.2	3.3	4.4	2.4	4.1	2.3	2.2	3.1	2.5	
POM Bext	9.6	10.9	5.7	6.2	7.1	14.8	10.4	5.6	14.7	6.1	12.1	16.0	
EC Bext	1.8	2.1	0.5	1.6	1.5	2.3	1.8	1.8	2.8	1.6	1.9	2.4	
Soil Bext	0.6	0.5	0.6	0.9	0.4	0.4	0.5	0.4	0.7	0.4	0.6	0.5	
Coarse Mass Bext	1.7	2.5	1.9	1.6	2.0	2.0	1.4	2.2	2.4	1.6	2.0	4.5	
Sea Salt Bext	0.4	0.1	0.5	0.7	0.5	0.3	0.1	0.3	0.0	0.1	0.2	0.1	




White Pass - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	1.7	1.8	0.8	2.2	1.9	1.1	1.5	1.9	1.8	1.8	1.5	1.7	
Sulfate Bext	0.9	1.1	0.2	1.4	1.0	0.5	0.6	1.4	0.9	1.1	0.9	1.0	
Nitrate Bext	0.2	0.2	0.2	0.4	0.3	0.1	0.2	0.3	0.1	0.2	0.1	0.2	
POM Bext	0.3	0.2	0.2	0.2	0.4	0.2	0.2	0.2	0.2	0.3	0.2	0.3	0.1
EC Bext	0.1	0.1	0.0	0.1	0.3	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1
Soil Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Coarse Mass Bext	0.1	0.2	0.1	0.2	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.4
Sea Salt Bext	0.1	0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.2	0.1	0.1	0.1




Mount Rainier - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	18.2	16.4	8.5	18.8	18.3	17.3		18.6	16.8	17.6	15.3	15.8	16.5
Sulfate Bext	23.7	17.3	2.2	25.6	28.6	18.6		22.1	14.9	21.0	15.0	15.8	19.6
Nitrate Bext	5.1	3.6	2.6	5.7	4.1	4.9		5.9	3.2	4.5	4.5	2.4	3.5
POM Bext	15.1	13.1	3.4	15.3	11.8	14.7		18.5	16.2	15.0	9.8	12.3	12.4
EC Bext	5.1	4.4	0.3	6.3	4.8	4.9		4.6	5.9	5.2	3.6	3.9	3.2
Soil Bext	0.5	0.5	0.6	0.3	0.6	0.6		0.6	0.4	0.6	0.4	0.6	0.4
Coarse Mass Bext	2.2	2.5	2.4	1.9	2.1	2.1		2.7	2.4	2.5	2.0	3.5	2.0
Sea Salt Bext	0.1	0.4	1.0	0.0	0.0	0.0		0.2	0.3	0.2	0.7	0.3	0.4




Mount Rainier - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	5.5	4.9	2.6	6.1	6.1	4.8		4.9	5.7	4.8	5.1	4.7	4.3
Sulfate Bext	2.6	2.5	0.3	3.1	3.0	2.1		2.1	3.2	2.0	2.3	2.5	2.2
Nitrate Bext	0.6	0.4	0.3	0.7	0.8	0.6		0.5	0.4	0.4	0.4	0.5	0.4
POM Bext	1.5	1.0	0.5	1.9	1.7	1.2		1.2	1.4	1.0	1.2	0.9	0.7
EC Bext	0.7	0.5	0.1	0.7	0.9	0.5		0.5	0.7	0.7	0.5	0.3	0.4
Soil Bext	0.0	0.0	0.1	0.1	0.1	0.0		0.0	0.0	0.0	0.0	0.0	0.1
Coarse Mass Bext	0.5	0.4	0.5	0.5	0.6	0.6		0.5	0.6	0.3	0.5	0.4	0.4
Sea Salt Bext	0.5	0.6	0.1	0.5	0.5	0.4		0.6	0.6	0.8	0.6	0.5	0.4




North Cascades - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	16.0	13.7	8.4	13.8	14.1	20.5	15.7	12.9	14.4	12.6	12.9	15.8	
Sulfate Bext	14.9	13.7	2.0	15.6	14.9	13.6	15.4	12.9	14.7	13.5	12.4	15.2	
Nitrate Bext	2.7	2.3	2.1	2.1	2.8	3.7	2.2	2.2	2.5	2.2	2.4	2.1	
POM Bext	33.0	9.4	6.5	7.6	9.0	98.7	16.8	6.4	9.4	5.9	7.3	18.1	
EC Bext	3.8	2.1	0.5	1.8	2.2	8.9	2.3	2.1	2.5	1.7	1.5	2.6	
Soil Bext	0.5	0.4	0.5	0.7	0.4	0.3	0.5	0.3	0.5	0.3	0.5	0.3	
Coarse Mass Bext	1.8	1.5	1.9	1.9	1.6	1.5	2.0	1.5	1.8	1.3	1.6	1.3	
Sea Salt Bext	0.0	0.2	0.2	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.3	0.1	




North Cascades - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	3.4	3.2	1.9	3.8	3.1	3.2	3.4	3.3	2.8	3.0	3.7	3.2	
Sulfate Bext	1.4	1.6	0.3	1.8	1.3	1.2	1.2	1.6	1.2	1.2	1.4	2.2	1.6
Nitrate Bext	0.4	0.2	0.3	0.5	0.4	0.4	0.5	0.2	0.2	0.2	0.2	0.3	0.2
POM Bext	0.6	0.5	0.3	0.7	0.5	0.7	0.7	0.5	0.3	0.3	0.5	0.5	0.5
EC Bext	0.2	0.2	0.0	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.1	0.1	0.2
Soil Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Coarse Mass Bext	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.1
Sea Salt Bext	0.1	0.1	0.0	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.1	0.1	0.1




Olympic - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	16.7	15.2	8.4	16.5	17.2	16.6	15.4	15.2	14.8	15.2	15.4		
Sulfate Bext	16.7	15.8	1.5	16.9	16.7	16.4	16.3	15.6	13.7	14.5	19.1		
Nitrate Bext	8.3	6.2	2.0	9.3	8.2	7.4	5.7	6.5	7.2	7.2	4.6		
POM Bext	12.1	7.8	3.1	10.9	14.0	11.2	8.8	7.5	7.0	8.3	7.5		
EC Bext	2.7	2.1	0.3	2.6	3.1	2.5	2.6	2.2	2.3	1.8	1.4		
Soil Bext	0.3	0.3	0.3	0.4	0.2	0.3	0.2	0.3	0.2	0.4	0.3		
Coarse Mass Bext	1.8	1.6	1.9	1.7	1.7	1.9	1.4	2.1	1.4	1.5	1.6		
Sea Salt Bext	1.4	1.4	3.4	0.3	1.6	2.5	1.0	1.1	1.2	1.4	2.0		




Olympic - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	6.0	5.0	2.7			5.5	6.0	6.5	5.9	5.2	5.1	5.0	4.1
Sulfate Bext	2.7	2.3	0.4			2.6	2.6	2.9	2.8	2.0	2.1	2.6	2.0
Nitrate Bext	1.2	0.8	0.3			1.0	1.4	1.4	1.1	0.8	0.9	0.7	0.7
POM Bext	1.9	1.1	0.7			1.6	1.9	2.2	1.4	1.1	1.3	1.2	0.6
EC Bext	0.6	0.4	0.1			0.6	0.6	0.8	0.5	0.5	0.5	0.4	0.2
Soil Bext	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Coarse Mass Bext	0.3	0.3	0.3			0.4	0.3	0.3	0.3	0.3	0.2	0.3	0.2
Sea Salt Bext	0.6	0.8	0.3			0.4	0.6	0.9	1.2	1.2	0.6	0.5	0.5




Pasayten - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	15.2	14.1	8.3	15.7	14.1	15.4	15.7	12.4	19.4	13.8	12.2	12.7	
Sulfate Bext	8.1	9.5	1.7	9.7	7.1	7.9	7.5	10.9	8.1	10.3	9.3	8.8	
Nitrate Bext	3.3	3.2	2.2	4.8	3.5	3.5	1.3	4.7	2.8	3.3	3.8	1.3	
POM Bext	21.9	19.4	5.9	23.3	16.9	20.3	27.1	6.2	55.1	13.9	7.9	13.8	
EC Bext	3.3	3.0	0.6	3.9	3.0	3.2	3.2	1.8	7.5	2.5	1.4	1.9	
Soil Bext	0.8	0.6	0.8	1.4	0.7	0.6	0.5	0.5	0.8	0.6	0.7	0.5	
Coarse Mass Bext	2.1	2.1	2.2	2.6	1.6	2.2	1.9	1.3	2.8	2.2	1.9	2.2	
Sea Salt Bext	0.1	0.1	0.2	0.2	0.0	0.1	0.0	0.2	0.0	0.1	0.1	0.2	



Pasayten - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.7	2.5	1.2		3.0	2.6	2.5	2.9	3.2	2.0	2.4	2.7	2.3
Sulfate Bext	1.5	1.5	0.4		1.7	1.4	1.4	1.6	1.9	1.1	1.4	1.9	1.4
Nitrate Bext	0.5	0.4	0.3		0.6	0.6	0.4	0.5	0.5	0.3	0.5	0.4	0.5
POM Bext	0.5	0.4	0.2		0.6	0.5	0.6	0.6	0.6	0.4	0.3	0.4	0.2
EC Bext	0.2	0.1	0.0		0.2	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.1
Soil Bext	0.1	0.1	0.1		0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1
Coarse Mass Bext	0.2	0.2	0.1		0.1	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
Sea Salt Bext	0.1	0.2	0.1		0.1	0.2	0.1	0.1	0.3	0.1	0.3	0.1	0.1




Dolly Sods - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000	2005	2064	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	29.0	27.6	10.4	29.0	28.2	28.5	29.7	29.8	30.9	29.8	29.5	25.4	22.2
Sulfate Bext	160.5	141.3	3.2	148.6	155.1	153.3	171.2	174.4	195.3	176.9	164.3	100.4	69.7
Nitrate Bext	3.4	2.0	1.2	3.8	4.1	2.9	4.4	1.7	1.7	1.8	2.3	2.5	1.6
POM Bext	13.4	11.5	10.5	15.4	11.5	11.8	13.5	14.6	10.2	11.8	15.7	12.6	7.3
EC Bext	4.7	3.7	0.3	6.4	4.2	4.1	4.9	3.7	4.4	4.3	4.0	3.2	2.3
Soil Bext	0.8	0.8	0.8	0.6	0.8	0.8	0.6	1.1	0.7	0.7	0.9	0.9	0.7
Coarse Mass Bext	1.7	2.0	2.3	2.8	1.8	1.6	1.0	1.5	2.1	1.5	2.6	2.2	1.6
Sea Salt Bext	0.1	0.1	0.3	0.0	0.3	0.0	0.0	0.3	0.1	0.2	0.1	0.1	0.0




Dolly Sods - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	12.3	10.2	3.6	13.0	13.3	11.9	11.5	11.7	12.1	10.6	10.3	9.4	8.7
Sulfate Bext	14.3	11.2	0.8	14.1	18.0	13.2	13.0	13.4	14.2	12.0	11.6	9.7	8.3
Nitrate Bext	3.1	2.1	0.4	4.0	2.9	3.0	3.1	2.5	3.5	1.6	1.6	2.0	1.6
POM Bext	4.9	2.9	2.4	6.6	4.7	4.6	4.4	4.0	3.6	2.9	3.1	2.4	2.5
EC Bext	2.1	1.4	0.1	2.9	2.1	2.0	1.7	1.6	1.9	1.6	1.4	1.0	1.0
Soil Bext	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.3
Coarse Mass Bext	0.7	0.8	0.6	0.7	0.7	0.7	0.7	0.8	0.9	0.8	0.8	0.8	0.7
Sea Salt Bext	0.1	0.2	0.1	0.1	0.0	0.0	0.1	0.4	0.1	0.2	0.2	0.3	0.1




Bridger - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.1	10.7	6.5	11.0	10.9	12.1	11.4	10.2	9.1	11.9	11.0	12.5	8.8
Sulfate Bext	5.0	5.2	0.8	4.5	5.9	4.3	4.9	5.3	6.3	5.0	5.1	5.4	4.1
Nitrate Bext	1.4	1.1	0.8	1.3	1.5	1.7	1.1	1.6	0.8	1.2	1.4	1.4	0.9
POM Bext	10.5	11.1	4.6	10.3	7.6	13.8	14.1	6.9	5.0	13.1	10.0	20.3	7.3
EC Bext	2.0	1.9	0.4	2.6	1.5	2.4	2.1	1.4	1.3	2.5	1.9	2.5	1.3
Soil Bext	1.1	1.0	1.1	0.9	1.5	1.1	0.7	1.3	0.8	1.1	0.9	1.4	0.9
Coarse Mass Bext	2.5	2.3	2.7	2.5	3.0	2.2	2.1	2.7	2.1	2.3	2.5	2.6	1.8
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.1	0.1




Bridger - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.1	1.5	0.3	2.5	2.5	2.1	1.6	1.8	1.6	1.5	2.0	1.3	1.3
Sulfate Bext	1.5	1.2	0.3	1.9	1.6	1.4	1.1	1.3	1.3	1.1	1.3	1.1	1.2
Nitrate Bext	0.4	0.3	0.2	0.4	0.5	0.5	0.3	0.4	0.2	0.3	0.4	0.4	0.3
POM Bext	0.8	0.6	0.4	1.0	0.7	0.7	0.8	0.7	0.7	0.6	0.8	0.5	0.4
EC Bext	0.4	0.2	0.1	0.3	0.6	0.4	0.3	0.2	0.3	0.2	0.3	0.1	0.1
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.2
Sea Salt Bext	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0




North Absaroka - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.5	11.0	6.8		10.5	13.8	10.1	9.6	11.6			11.8	
Sulfate Bext	4.9	4.6	0.8		5.3	4.7	4.6	4.8	4.8			4.2	
Nitrate Bext	1.6	1.4	0.8		2.3	1.1	1.5	1.4	1.1			1.5	
POM Bext	11.6	10.0	4.6		6.5	20.1	8.4	6.0	10.5			13.5	
EC Bext	1.9	1.8	0.4		1.3	2.9	1.4	1.6	2.1			1.7	
Soil Bext	0.8	0.8	0.9		1.0	0.8	0.7	0.6	1.2			0.7	
Coarse Mass Bext	2.9	3.4	3.4		3.4	2.9	2.4	2.9	4.4			2.9	
Sea Salt Bext	0.0	0.0	0.0		0.0	0.0	0.0	0.0	0.0			0.0	




North Absaroka - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.0	1.2	0.6			2.2	1.9	2.0	1.3	1.1		1.3	
Sulfate Bext	1.1	1.0	0.3			1.3	0.9	1.1	1.0	0.9		1.2	
Nitrate Bext	0.4	0.2	0.2			0.5	0.3	0.4	0.2	0.2		0.3	
POM Bext	0.8	0.4	0.5			0.8	0.8	0.9	0.5	0.3		0.3	
EC Bext	0.2	0.2	0.0			0.2	0.2	0.2	0.2	0.2		0.1	
Soil Bext	0.1	0.1	0.1			0.1	0.1	0.1	0.1	0.1		0.1	
Coarse Mass Bext	0.7	0.4	0.5			0.7	0.8	0.6	0.4	0.5		0.4	
Sea Salt Bext	0.0	0.0	0.0			0.0	0.0	0.0	0.0	0.0		0.0	



Yellowstone - Worst 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	11.8	11.5	6.4	12.9	12.6	11.4	11.2	10.8	9.6	11.8	12.6	14.0	9.6
Sulfate Bext	4.3	4.6	0.8	4.2	4.7	3.8	4.1	4.6	4.8	5.1	4.3	5.0	3.5
Nitrate Bext	1.8	1.6	0.6	2.0	2.4	1.6	1.4	1.5	1.6	0.9	2.4	1.7	1.6
POM Bext	13.5	15.5	4.6	18.4	16.1	11.2	12.8	8.9	6.7	15.1	14.7	25.6	15.2
EC Bext	2.5	2.2	0.4	3.7	2.6	1.9	2.3	1.9	1.8	2.3	2.4	2.6	1.7
Soil Bext	1.0	0.8	1.0	0.8	1.4	1.2	0.4	0.9	0.6	1.1	0.8	1.0	0.6
Coarse Mass Bext	2.6	2.4	3.0	2.8	2.7	3.1	1.3	2.9	1.8	2.6	2.6	2.6	2.3
Sea Salt Bext	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.1	0.1




Yellowstone - Best 20% Visibility Days – 5-Year Average, Natural Conditions and Annual Data

Parameter	2000-2004	2005-2009	2064 NC	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Deciview (dv)	2.6	2.0	0.4	2.8	3.0	2.6	1.9	2.6	2.1	2.3	2.2	1.8	1.9
Sulfate Bext	1.5	1.4	0.3	1.7	1.9	1.3	1.1	1.3	1.5	1.6	1.2	1.1	1.4
Nitrate Bext	0.7	0.5	0.3	0.9	0.8	0.8	0.5	0.6	0.3	0.4	0.7	0.6	0.6
POM Bext	1.1	0.8	0.5	1.1	1.2	1.1	0.9	1.3	1.1	0.8	1.0	0.8	0.5
EC Bext	0.3	0.3	0.1	0.2	0.4	0.3	0.2	0.5	0.3	0.4	0.3	0.1	0.2
Soil Bext	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Coarse Mass Bext	0.2	0.2	0.2	0.3	0.2	0.3	0.2	0.3	0.1	0.2	0.2	0.3	0.3
Sea Salt Bext	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

APPENDIX G.2. SPATIAL MAPS OF REGIONAL HAZE RULE METRICS

G.2.1. Comparisons of progress to the URP goal

The change in the 20% worst haze RHR metric can be examined two ways: as the percent change and as the absolute change compared to the URP goal for each site (i.e., the difference between period 1 and baseline divided by the deciview change required to attain the URP for each site). Color dot maps depicting the fractional and absolute worst haze RHR metric changes are provided here.



Fraction of dv uniform rate of progress (URP) from the baseline (2000–2004) to period 1 (2005–2009) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites. Brown circles indicate degradation in the worst 20% visibility days, while blue circles represent improvement in worst 20% visibility days. The two darkest shades of blue indicate progress that is at or better than the URP.

Tracking Progress 2000-2004 to 2005-2009
 Hazeiest 20% Days: Period 1 - Baseline (dv)



Absolute change in dv from the baseline (2000–2004) to period 1 (2005–2009) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites. Brown circles indicate degradation in the worst 20% visibility days, while blue circles represent improvement in worst 20% visibility days.

Tracking Progress 2000-2004 to 2005-2009
 Clearest 20% Days: Period 1 - Baseline (dv)




Absolute change in dv from the baseline (2000–2004) to period 1 (2005–2009) for the 20% best visibility days at 107 of the 110 IMPROVE regional haze tracking sites. Brown circles indicate degradation in the best 20% visibility days, while blue circles represent improvement in best 20% visibility days.



Absolute change in annual mean dv from the baseline (2000–2004) to period 1 (2005–2009) at 107 of the 110 IMPROVE regional haze tracking sites. Brown circles indicate degradation in annual mean dv , while blue circles represent improvement in annual mean dv .

G.2.2 Fraction of hypothetical uniform rate of progress (URP) for individual species

While there are no official URP specified in the RHR for any of the aerosol species b_{ext} components, hypothetical URP values were generated for each site by reducing each component of b_{ext} by the same fraction that would be required in total extinction to evaluate progress toward the conceptual URP in dv .



Fraction of hypothetical ammonium sulfate uniform rate of progress (URP) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate degradation in the worst 20% visibility days due to ammonium sulfate extinction, while blue circles represent improvement in worst 20% visibility days due to ammonium sulfate extinction. Only the two darkest blue colored circles indicate progress that is at or better than the hypothetical ammonium sulfate extinction URP.




Fraction of hypothetical ammonium nitrate uniform rate of progress (URP) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate degradation in the worst 20% visibility days due to ammonium nitrate extinction, while blue circles represent improvement in worst 20% visibility days due to ammonium nitrate extinction.

extinction. Only the two darkest blue colored circles indicate progress that is at or better than the hypothetical ammonium nitrate extinction URP.



Fraction of hypothetical particulate organic matter uniform rate of progress (URP) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate degradation in the worst 20% visibility days due to particulate organic matter extinction, while blue circles represent improvement in worst 20% visibility days due to particulate organic matter extinction. Only the two darkest blue colored circles indicate progress that is at or better than the hypothetical particulate organic matter extinction URP.




Fraction of hypothetical elemental carbon uniform rate of progress (URP) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate degradation in the worst 20% visibility days due to elemental

carbon extinction, while blue circles represent improvement in worst 20% visibility days due to elemental carbon extinction. Only the two darkest blue colored circles indicate progress that is at or better than the hypothetical elemental carbon extinction URP.



Fraction of hypothetical soil uniform rate of progress (URP) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate degradation in the worst 20% visibility days due to soil extinction, while blue circles represent improvement in worst 20% visibility days due to soil extinction. Only the two darkest blue colored circles indicate progress that is at or better than the hypothetical soil extinction URP.



Fraction of hypothetical coarse mass uniform rate of progress (URP) for the 20% worst visibility days at 107 of the 110 IMPROVE regional haze tracking sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate degradation in the worst 20% visibility days due to coarse mass extinction, while blue circles represent improvement in worst 20% visibility days due to coarse mass extinction. Only the two

darkest blue colored circles indicate progress that is at or better than the hypothetical coarse mass extinction URP.

G.2.3 Absolute Difference in light extinction coefficients for individual species



Change in 20% worst day ammonium sulfate extinction ($b_{\text{ext_AS}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_AS}}$ on haziest 20% days, while blue circles represent decreases in $b_{\text{ext_AS}}$.



Change in 20% worst day ammonium nitrate extinction ($b_{\text{ext_AN}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_AN}}$ on haziest 20% days, while blue circles represent decreases in $b_{\text{ext_AN}}$.

Tracking Progress 2000-2004 to 2005-2009
 Haziest 20% Days: Period1-Baseline (Organic Bext Mm-1)



Change in 20% worst day particulate organic matter extinction ($b_{\text{ext_POM}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_POM}}$ on haziest 20% days, while blue circles represent decreases in $b_{\text{ext_POM}}$.

Tracking Progress 2000-2004 to 2005-2009
 Haziest 20% Days: Period1-Baseline (EC Bext Mm-1)



Change in 20% worst day elemental carbon extinction ($b_{\text{ext_EC}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_EC}}$ on haziest 20% days, while blue circles represent decreases in $b_{\text{ext_EC}}$.

Tracking Progress 2000-2004 to 2005-2009
 Haziest 20% Days: Period1-Baseline (Fine Soil Bext Mm⁻¹)



Change in 20% worst day soil extinction ($b_{\text{ext_soil}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_soil}}$ on haziest 20% days, while blue circles represent decreases in $b_{\text{ext_soil}}$.

Tracking Progress 2000-2004 to 2005-2009
 Haziest 20% Days: Period1-Baseline (Coarse Mass Bext Mm⁻¹)



Change in 20% worst day coarse mass extinction ($b_{\text{ext_CM}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_CM}}$ on haziest 20% days, while blue circles represent decreases in $b_{\text{ext_CM}}$.

Tracking Progress 2000-2004 to 2005-2009

Haziest 20% Days: Period1-Baseline (Sea Salt Bext Mm⁻¹)



Change in 20% worst day sea salt extinction (b_{ext_SS} , Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in b_{ext_SS} on haziest 20% days, while blue circles represent decreases in b_{ext_SS} .

Tracking Progress 2000-2004 to 2005-2009

Annual Mean: Period1-Baseline (Sulfate Bext Mm⁻¹)



Change in annual mean ammonium sulfate extinction (b_{ext_AS} , Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean b_{ext_AS} , while blue circles represent decreases in b_{ext_AS} .

Tracking Progress 2000-2004 to 2005-2009
Annual Mean: Period1-Baseline (Nitrate Bext Mm⁻¹)



Change in annual mean ammonium nitrate extinction ($b_{\text{ext_AN}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean $b_{\text{ext_AN}}$, while blue circles represent decreases in $b_{\text{ext_AN}}$.

Tracking Progress 2000-2004 to 2005-2009
Annual Mean: Period1-Baseline (Organic Bext Mm⁻¹)



Change in annual mean particulate organic matter extinction ($b_{\text{ext_POM}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean $b_{\text{ext_POM}}$, while blue circles represent decreases in $b_{\text{ext_POM}}$.

Tracking Progress 2000-2004 to 2005-2009
Annual Mean: Period1-Baseline (EC Bext Mm-1)



Change in annual mean elemental carbon extinction ($b_{\text{ext_EC}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean $b_{\text{ext_EC}}$, while blue circles represent decreases in $b_{\text{ext_EC}}$.

Tracking Progress 2000-2004 to 2005-2009
Annual Mean: Period1-Baseline (Fine Soil Bext Mm-1)



Change in annual mean soil extinction ($b_{\text{ext_soil}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean $b_{\text{ext_soil}}$, while blue circles represent decreases in $b_{\text{ext_soil}}$.

Tracking Progress 2000-2004 to 2005-2009

Annual Mean: Period1-Baseline (Coarse Mass Bext Mm⁻¹)



Change in annual mean coarse mass extinction ($b_{\text{ext_CM}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean $b_{\text{ext_CM}}$, while blue circles represent decreases in $b_{\text{ext_CM}}$.

Tracking Progress 2000-2004 to 2005-2009

Annual Mean: Period1-Baseline (Sea Salt Bext Mm⁻¹)



Change in annual mean sea salt extinction ($b_{\text{ext_SS}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in annual mean $b_{\text{ext_SS}}$, while blue circles represent decreases in $b_{\text{ext_SS}}$.

Tracking Progress 2000-2004 to 2005-2009
 Clearest 20% Days: Period1-Baseline (Sulfate Bext Mm⁻¹)



Change in 20% best day ammonium sulfate extinction ($b_{\text{ext_AS}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_AS}}$ on best 20% days, while blue circles represent decreases in $b_{\text{ext_AS}}$.

Tracking Progress 2000-2004 to 2005-2009
 Clearest 20% Days: Period1-Baseline (Nitrate Bext Mm⁻¹)



Change in 20% best day ammonium nitrate extinction ($b_{\text{ext_AN}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_AN}}$ on best 20% days, while blue circles represent decreases in $b_{\text{ext_AN}}$.

Tracking Progress 2000-2004 to 2005-2009

Clearest 20% Days: Period1-Baseline (Organic Bext Mm-1)



Change in 20% best day particulate organic matter extinction ($b_{\text{ext_POM}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_POM}}$ on best 20% days, while blue circles represent decreases in $b_{\text{ext_POM}}$.

Tracking Progress 2000-2004 to 2005-2009

Clearest 20% Days: Period1-Baseline (EC Bext Mm-1)



Change in 20% best day elemental carbon extinction ($b_{\text{ext_EC}}$, Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_EC}}$ on best 20% days, while blue circles represent decreases in $b_{\text{ext_EC}}$.

Tracking Progress 2000-2004 to 2005-2009
 Clearest 20% Days: Period1-Baseline (Fine Soil Bext Mm-1)



Change in 20% best day soil extinction ($b_{\text{ext_soil}}, \text{Mm}^{-1}$) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_soil}}$ on best 20% days, while blue circles represent decreases in $b_{\text{ext_soil}}$.

Tracking Progress 2000-2004 to 2005-2009
 Clearest 20% Days: Period1-Baseline (Coarse Mass Bext Mm-1)



Change in 20% best day coarse mass extinction ($b_{\text{ext_CM}}, \text{Mm}^{-1}$) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in $b_{\text{ext_CM}}$ on best 20% days, while blue circles represent decreases in $b_{\text{ext_CM}}$.

Tracking Progress 2000-2004 to 2005-2009
Clearest 20% Days: Period1-Baseline (Sea Salt Bext Mm⁻¹)



Change in 20% best day sea salt extinction (b_{ext_SS} , Mm^{-1}) at 107 of the 110 IMPROVE sites from the baseline (2000–2004) to period 1 (2005–2009). Brown circles indicate increases in b_{ext_SS} on best 20% days, while blue circles represent decreases in b_{ext_SS} .