

Contents

5	Light Extinction Budget	5-1
5.1	Assumptions and Data Analysis Techniques	5-1
5.1.1	Available Data	5-1
5.1.2	Chemical Composition of Particulate Matter	5-1
5.1.3	Relative Humidity Considerations	5-2
5.1.4	Multiple Linear Regression Analysis	5-2
5.1.5	Consensus Scattering and Absorption Efficiencies	5-7
5.1.6	Reconstructed Scattering and Extinction	5-7
5.2	Preliminary Data Analysis	5-8
5.2.1	Optical Data and Relative Humidity Effects	5-9
5.2.2	Examination of Carbon Data	5-14
5.3	Light Extinction Budget By Extinction Type	5-24
5.3.1	Scattering by gases (Rayleigh Scattering)	5-24
5.3.2	Absorption by gases (NO ₂ Absorption)	5-30
5.3.3	Coarse Particle Scattering	5-32
5.3.4	Particle Absorption	5-32
5.3.5	Fine Particle Scattering	5-32
5.4	Extinction Budget by Species	5-36
5.4.1	Sulfates	5-36
5.4.2	Organics	5-37
5.4.3	Nitrates	5-40
5.4.4	Fine Soil Dust	5-40
5.5	Time Variations in the Light Extinction Budget	5-40
5.5.1	Extinction budgets for Individual Time Periods	5-40
5.5.2	Budgets on Extreme Extinction Days	5-58
5.6	Comparison to Previous Studies at Grand Canyon	5-58
5.7	Summary	5-58

List of Tables

5.1	Pearson correlation coefficients for data at Page.	5-4
5.2	Pearson correlation coefficients for Canyonlands.	5-4
5.3	Pearson correlation coefficients for data at Hopi Point.	5-4
5.4	Pearson correlation coefficients for Page, Canyonlands, and Hopi Point combined.	5-5
5.5	Rotated factor pattern for Page.	5-5
5.6	Rotated factor pattern for Canyonlands	5-5
5.7	Rotated factor pattern for Hopi Point	5-6
5.8	Rotated factor pattern for Page, Canyonlands, and Hopi Point combined.	5-6
5.9	Consensus literature light extinction efficiencies in m^2/g	5-7
5.10	Statistics for the 12-hour averaged optical data at Page (Km^{-1}). The low relative humidity subgroup is defined as $RH < 60\%$	5-15
5.11	Statistics for the 12-hour averaged optical data at Canyonlands (Km^{-1}). The low relative humidity subgroup is defined as $RH < 60\%$	5-15
5.12	Statistics for the 12-hour averaged optical data at Hopi Point (Km^{-1}). The low relative humidity subgroup is defined as $RH < 60\%$	5-16
5.13	Correlation matrix for carbonaceous material at Page.	5-16
5.14	Correlation matrix for carbonaceous material at Hopi Point.	5-16
5.15	Summary of multiple linear regression analyses using TOR and TMO Carbon. Results are the predicted light absorption efficiencies for absorbing carbon \pm the standard error in m^2/g . The dependent variable for all analyses was b_{ext} . Low RH refers to $RH < 60\%$	5-23
5.16	Statistics for the 12-hour averaged particulate and NO_2 data at Page, ($\mu g/m^3$). All data except coarse mass and NO_2 are for particles $< 2.5 \mu m$ diameter. Coarse mass is $2.5-15 \mu m$. Low RH is $RH < 60\%$	5-28
5.17	Statistics for the 12-hour averaged particulate data at Canyonlands, ($\mu g/m^3$). All data are for particles $< 2.5 \mu m$ diameter. Low RH is $RH < 60\%$	5-28
5.18	Statistics for the 12-hour averaged particulate data at Hopi Point, ($\mu g/m^3$). All data are for particles $< 2.5 \mu m$ diameter. Coarse mass is $2.5-15 \mu m$. Low RH is $RH < 60\%$	5-30
5.19	Mean extinction budgets by type for Page, Canyonlands, and Hopi Point for the 6-week WHITEX study period.	5-31
5.20	Summary of multiple linear regression analyses for coarse mass. Results are scattering efficiencies in m^2/g	5-33
5.21	Summary of multiple linear regression analyses for fine mass. Results are scattering efficiencies in m^2/g . Low RH is $RH < 60\%$. (fine-abs. C) is the fine mass remaining after absorbing carbon has been subtracted.	5-35
5.22	Scattering coefficients (Km^{-1}) for fine mass for each site as estimated by 4 different methods	5-36

5.23	Average fine particle scattering due to each chemical species for Page, Canyonlands, and Hopi Point using both MLR derived and consensus literature values for the scattering efficiencies.	5-37
5.24	Summary of multiple linear regression analyses used to determine sulfate scattering efficiency for 0% RH. Results are in m^2/g	5-38
5.25	Summary of multiple linear regression analyses used to determine the scattering efficiency for organics. Results are in m^2/g	5-39
5.26	Summary of multiple linear regression analyses used to determine nitrate scattering efficiency for 0% RH. Results are in m^2/g	5-41
5.27	Summary of multiple linear regression analyses used to determine fine soil scattering efficiency. Results are in m^2/g	5-42
5.28	Extinction coefficients (Km^{-1}) due to each chemical species at Page. The total and measured extinction do not include Rayleigh scattering.	5-43
5.28	Extinction coefficients (Km^{-1}) due to each chemical species at Page. The total and measured extinction do not include Rayleigh scattering (continued from previous page).	5-44
5.29	Fraction of non-Rayleigh extinction due to each chemical species at Page.	5-45
5.29	Fraction of non-Rayleigh extinction due to each chemical species at Page (continued from previous page).	5-46
5.30	Extinction coefficients (Km^{-1}) due to each chemical species at Canyonlands. The total and measured extinction do not include Rayleigh scattering.	5-47
5.30	Extinction coefficients (Km^{-1}) due to each chemical species at Canyonlands. The total and measured extinction do not include Rayleigh scattering (continued from previous page).	5-48
5.31	Fraction of non-Rayleigh extinction due to each chemical species at Canyonlands.	5-49
5.31	Fraction of non-Rayleigh extinction due to each chemical species at Canyonlands (continued from previous page).	5-50
5.32	Extinction coefficients (Km^{-1}) due to each chemical species at Hopi Point when maximum organics are used. The total and measured extinction do not include Rayleigh scattering.	5-51
5.32	Extinction coefficients (Km^{-1}) due to each chemical species at Hopi Point when maximum organics are used. The total and measured extinction do not include Rayleigh scattering (continued from previous page).	5-52
5.33	Fraction of non-Rayleigh light extinction due to each chemical species at Hopi Point when maximum organics are used.	5-53
5.33	Fraction of non-Rayleigh light extinction due to each chemical species at Hopi Point when maximum organics are used (continued from previous page).	5-54
5.34	Extinction coefficients (Km^{-1}) due to each chemical species at Hopi Point when minimum organics are used. The total and measured extinction do not include Rayleigh scattering.	5-55
5.34	Extinction coefficients (Km^{-1}) due to each chemical species at Hopi Point when minimum organics are used. The total and measured extinction do not include Rayleigh scattering (continued from previous page).	5-56
5.35	Fraction of non-Rayleigh extinction due to each chemical species at Hopi Point when minimum organics are used.	5-57
5.35	Fraction of non-Rayleigh extinction due to each chemical species at Hopi Point when minimum organics are used (continued from previous page).	5-58

5.36 24-hour average extinction budgets for extreme extinction days. Feb 12 is Julian day

43. Feb 14 is Julian day 45. 5-64

List of Figures

5.1	Relative light scattering coefficients for ammonium sulfate and ammonium nitrate aerosols.	5-3
5.2	Time traces of optical data and relative humidity at Page	5-10
5.3	Time traces of optical data and relative humidity at Canyonlands.	5-11
5.4	Time traces of optical data and relative humidity at Hopi Point.	5-12
5.5	Ratio of b_{ext}/b_{scat} as a function of relative humidity for Page.	5-13
5.6	Ratio of b_{ext}/b_{scat} as a function of relative humidity for Canyonlands.	5-13
5.7	Ratio of b_{ext}/b_{scat} as a function of relative humidity for Hopi Point.	5-14
5.8	Scatter plot of measured extinction vs reconstructed extinction for Page using TMO carbon. Sulfate and nitrate scattering efficiencies are both $2.55/(1-RH)$	5-18
5.9	Scatter plot of measured extinction vs reconstructed extinction for Page using TMO carbon. Sulfate scattering efficiency is $2.55 \times fs(RH)$. Nitrate scattering efficiency is $1.1 \times fn(RH)$	5-18
5.10	Scatter plot of measured extinction vs reconstructed extinction for Page using TOR carbon. Sulfate and nitrate scattering efficiencies are both $2.55/(1-RH)$	5-19
5.11	Scatter plot of measured extinction vs reconstructed extinction for Page using TOR carbon. Sulfate scattering efficiency is $2.55 \times fs(RH)$. Nitrate scattering efficiency is $1.1 \times fn(RH)$	5-19
5.12	Scatter plot of measured extinction vs reconstructed extinction for Hopi Point using TMO carbon. Sulfate and nitrate scattering efficiencies are both $2.55/(1-RH)$	5-20
5.13	Scatter plot of measured extinction vs reconstructed extinction for Hopi Point using TMO carbon. Sulfate scattering efficiency is $2.55 \times fs(RH)$. Nitrate scattering efficiency is $1.1 \times fn(RH)$	5-20
5.14	Scatter plot of measured extinction vs reconstructed extinction for Hopi Point using minimum TOR carbon. Sulfate and nitrate scattering efficiencies are both $2.55/(1-RH)$	5-21
5.15	Scatter plot of measured extinction vs reconstructed extinction for Hopi Point using minimum TOR carbon. Sulfate scattering efficiency is $2.55 \times fs(RH)$. Nitrate scattering efficiency is $1.1 \times fn(RH)$	5-21
5.16	Scatter plot of measured extinction vs reconstructed extinction for Hopi Point using maximum TOR carbon. Sulfate and nitrate scattering efficiencies are both $2.55/(1-RH)$	5-22
5.17	Scatter plot of measured extinction vs reconstructed extinction for Hopi Point using maximum TOR carbon. Sulfate scattering efficiency is $2.55 \times fs(RH)$. Nitrate scattering efficiency is $1.1 \times fn(RH)$	5-22
5.18	Time traces of particulate and NO_2 data ($\mu g/m^3$) at Page.	5-25
5.19	Time traces of particulate data ($\mu g/m^3$) at Canyonlands.	5-26
5.20	Time traces of particulate data ($\mu g/m^3$) at Hopi Point.	5-27

5.21 Mean light extinction budgets.	5-29
5.22 Measured and reconstructed 12-hour averaged extinction coefficients (Km^{-1}) at Page. Rayleigh scattering is included.	5-60
5.23 Measured and reconstructed 12-hour averaged extinction coefficients (Km^{-1}) at Canyonlands. Rayleigh scattering is included.	5-61
5.24 Measured and reconstructed 12-hour averaged extinction coefficients (Km^{-1}) at Hopi Point when minimum organics are used. Rayleigh scattering is included.	5-62
5.25 Measured and reconstructed 12-hour averaged extinction coefficients (Km^{-1}) at Hopi Point when maximum organics are used. Rayleigh scattering is included.	5-63
5.26 Extinction budgets on extreme days at Page. Areas of the wedges are proportional to the light extinction for each component. Total reconstructed extinction is $0.0742 \pm 0.0070 \text{ Km}^{-1}$ on Feb 12 (Julian day 43) and $0.0116 \pm 0.0014 \text{ Km}^{-1}$ on Feb 14 (Julian day 45).	5-65
5.27 Extinction budgets on Feb 12 and 14 at Canyonlands. Areas of the wedges are proportional to the light extinction for each component. Total reconstructed extinction is $0.0716 \pm 0.0126 \text{ Km}^{-1}$ on Feb 12 (Julian day 43) and $0.0106 \pm 0.0015 \text{ Km}^{-1}$ on Feb 14 (Julian day 45).	5-66
5.28 Extinction budgets on Feb 11 and 14 at Hopi Point using minimum organics. Areas of the wedges are proportional to the light extinction for each component. Total reconstructed extinction is $0.0584 \pm 0.0138 \text{ Km}^{-1}$ on Feb 11 (Julian day 42) and $0.0097 \pm 0.0015 \text{ Km}^{-1}$ on Feb 14 (Julian day 45).	5-67
5.29 Extinction budgets on Feb 11 and 14 at Hopi Point using maximum organics. Areas of the wedges are proportional to the light extinction for each component. Total reconstructed extinction is $0.0604 \pm 0.0138 \text{ Km}^{-1}$ on Feb 11 (Julian day 42) and $0.0111 \pm 0.0015 \text{ Km}^{-1}$ on Feb 14 (Julian day 45).	5-68